

jc542 U.S. PTO



10/27/99

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

FIRST-NAMED INVENTOR: Murdin

FOR: *CHLAMYDIA* ANTIGENS AND CORRESPONDING DNA FRAGMENTS
AND USES THEREOF

October 27, 1999

Boston, Massachusetts

Box PATENT APPLICATION
Assistant Commissioner for Patents
Washington, D.C. 20231REQUEST FOR FILING A NEW NONPROVISIONAL APPLICATION
UNDER 37 C.F.R. §1.53(b)

1. This is a request for filing a new nonprovisional application under 37 C.F.R. §1.53(b).
2. ☒ Specification and Drawings (Total pages: 87);
Specification (62 pages); Claims (5 pages); Abstract (1 page);
and Drawings: 19 sheets; FIGS. 1-4.
☒ Formal
☐ Informal
3. ☒ Declaration and Power of Attorney
☒ Unsigned
☐ Signed
4. ☐ Information Disclosure Statement (IDS)
☐ Copy of IDS and PTO-1449 (___ pages)
☐ Copies of references cited
5. ☐ Assignment Papers
☐ Recordation Form Cover Sheet (PTO-1595)
☐ Assignment Document
6. ☐ Statement Claiming Small Entity Status
☐ Claiming Small Entity As Independent Inventor (37 C.F.R. §§1.9(f) & 1.27(b)).
☐ Claiming Small Entity As Small Business Concern (37 C.F.R. §§1.9(f) & 1.27(c)).
☐ Claiming Small Entity As Nonprofit Organization (37 C.F.R. §§1.9(f) & 1.27(d)).

Applicant(s): Murdin, *et al.*


Request for New Nonprovisional Application (37 C.F.R. §1.53(b))

7. Fee Calculation

CLAIMS AS FILED					
Claims	Number Filed	Basic Fee Allowance	Number Extra	Rate	Basic Fee 37 C.F.R. 1.16(a) \$760.00
Total Claims (37 C.F.R. 1.16(c))	37	- 20 =	17	\$ 18.00	306.00
Independent Claims (37 C.F.R. 1.16(b))	4	- 3 =	1	\$78.00	78.00
Multiple Dependent Claim(s), if any (37 C.F.R. 1.16(d))	0			\$260.00	0
SUBTOTAL:					\$1,144.00
Reduction by 50% for filing by small entity:					- \$0
TOTAL FEE:					\$1,144.00

8. ☒ A check in the amount of **\$1,144.00** is enclosed.
9. ☒ The Commissioner is hereby authorized to credit overpayments or charge the following fees to Deposit Account No. 50-0311, Ref. No.19721-010:
- ☒ Fees required under 37 C.F.R. §1.16;
- ☒ Fees required under 37 C.F.R. §1.17;
- ☒ Fees required under 37 C.F.R. §1.18.
10. ☒ Return Receipt Postcard Enclosed.
11. ☒ Other Documents Enclosed:
- ☐ Change of Attorney Address In Application.
- ☒ Limited Recognition under 37 C.F.R. § 10.9(b) for Michel Morency.

Respectfully submitted,


Ivor R. Elrifi, Reg. No. 39,529

Michel Morency, Limited Recognition
Shelby J. Walker, Reg. No. 45,192
Attorneys for Applicants
MINTZ, LEVIN, COHN, FERRIS,
GLOVSKY and POPEO, P.C.
One Financial Center
Boston, Massachusetts 02111
Tel: (617) 542-6000
Fax: (617) 542-2241

Dated: October 27, 1999

CHLAMYDIA ANTIGENS AND CORRESPONDING DNA FRAGMENTS AND USES THEREOF

5 RELATED U.S. APPLICATION

The present patent application claims priority to the following United States provisional patent applications: U.S.S.Ns. 60/106,070, filed October 29, 1998 and 60/122,066, filed March 1, 1999, each incorporated herein by reference.

FIELD OF THE INVENTION

10 The present invention relates to *Chlamydia* antigens and corresponding DNA molecules, which can be used in methods to prevent and treat disease caused by *Chlamydia* infection in mammals, such as humans.

BACKGROUND OF THE INVENTION

Chlamydiae are prokaryotes. They exhibit morphologic and structural similarities to
15 Gram negative bacteria including a trilaminar outer membrane, which contains lipopolysaccharide and several membrane proteins. Chlamydiae are differentiated from other bacteria by their morphology and by a unique developmental cycle. They are obligate intracellular parasites with a unique biphasic life cycle consisting of a metabolically inactive but infectious extracellular stage and a replicating but non-infectious intracellular stage. The
20 replicative stage of the life-cycle takes place within a membrane-bound inclusion which sequesters the bacteria away from the cytoplasm of the infected host cell.

Because chlamydiae are small and multiply only within susceptible cells they were long thought to be viruses. However, they have many characteristics in common with other bacteria: (1) they contain both DNA and RNA, (2) they divide by binary fission, (3) their cell envelopes
25 resemble those of other Gram-negative bacteria, (4) they contain ribosomes similar to those of

other bacteria, and (5) they are susceptible to various antibiotics. Chlamydiae can be seen in the light microscope, and the genome is about one-third the size of the *Escherichia coli* genome.

Many different strains of chlamydiae have been isolated from birds, man, and other mammals, and these strains can be distinguished on the basis of host range, virulence,
5 pathogenesis, and antigenic composition. There is strong homology of DNA within each species, but surprisingly little between species, suggesting long-standing evolutionary separation.

C. trachomatis has a high degree of host specificity, being almost completely limited to man; it causes ocular and genitourinary infections of widely varying severity. In contrast, *C. psittaci* strains are rare in man but are found in a wide range of birds and also in wild,
10 domestic, and laboratory mammals, where they multiply in cells of many organs.

C. pneumoniae is a common human pathogen, originally described as the TWAR strain of *C. psittaci*, but subsequently recognized to be a new species. *C. pneumoniae* is antigenically, genetically, and morphologically distinct from other *Chlamydia* species (*C. trachomatis*, *C. pecorum* and *C. psittaci*). It shows 10% or less DNA sequence homology with either of
15 *C. trachomatis* or *C. psittaci* and so far appears to consist of only a single strain, TWAR.

C. pneumoniae is a common cause of community acquired pneumonia, less frequent only than *Streptococcus pneumoniae* and *Mycoplasma pneumoniae*. Grayston *et al.*, *J. Infect. Dis.* 168: 1231 (1995); Campos *et al.*, *Invest. Ophthalmol. Vis. Sci.* 36: 1477 (1995), each incorporated herein by reference. It can also cause upper respiratory tract symptoms and disease,
20 including bronchitis and sinusitis. See, *e.g.*, Grayston *et al.*, *J. Infect. Dis.* 168: 1231 (1995); Campos *et al.*, *Invest. Ophthalmol. Vis. Sci.* 36: 1477 (1995); Grayston *et al.*, *J. Infect. Dis.* 161: 618 (1990); Marrie, *Clin. Infect. Dis.* 18: 501 (1993). The great majority of the adult population (over 60%) has antibodies to *C. pneumoniae* (Wang *et al.*, *Chlamydial Infections*, Cambridge University Press, Cambridge, p. 329 (1986)), indicating past infection which was unrecognized
25 or asymptomatic.

C. pneumoniae infection usually presents as an acute respiratory disease (*i.e.*, cough, sore throat, hoarseness, and fever; abnormal chest sounds on auscultation). For most patients, the cough persists for 2 to 6 weeks, and recovery is slow. In approximately 10% of these cases,

upper respiratory tract infection is followed by bronchitis or pneumonia. Furthermore, during a *C. pneumoniae* epidemic, subsequent co-infection with pneumococcus has been noted in about half of these pneumonia patients, particularly in the infirm and the elderly. As noted above, there is more and more evidence that *C. pneumoniae* infection is also linked to diseases other than respiratory infections.

The reservoir for the organism is presumably people. In contrast to *C. psittaci* infections, there is no known bird or animal reservoir. Transmission has not been clearly defined. It may result from direct contact with secretions, from fomites, or from airborne spread. There is a long incubation period, which may last for many months. Based on analysis of epidemics, *C. pneumoniae* appears to spread slowly through a population (case-to-case interval averaging 30 days) because infected persons are inefficient transmitters of the organism. Susceptibility to *C. pneumoniae* is universal. Reinfections occur during adulthood, following the primary infection as a child. *C. pneumoniae* appears to be an endemic disease throughout the world, noteworthy for superimposed intervals of increased incidence (epidemics) that persist for 2 to 3 years. *C. trachomatis* infection does not confer cross-immunity to *C. pneumoniae*. Infections are easily treated with oral antibiotics, tetracycline or erythromycin (2 g/day, for at least 10 to 14 days). A recently developed drug, azithromycin, is highly effective as a single-dose therapy against chlamydial infections.

In most instances, *C. pneumoniae* infection is mild and without complications, and up to 90% of infections are subacute or unrecognized. Among children in industrialized countries, infections have been thought to be rare up to the age of five years, although a recent study has reported that many children in this age group show PCR evidence of infection despite being seronegative, and estimates a prevalence of 17-19% in 2-4 years old. See, Normann *et al.*, *Acta Paediatrica*, 87: 23-27 (1998). In developing countries, the seroprevalence of *C. pneumoniae* antibodies among young children is elevated, and there are suspicions that *C. pneumoniae* may be an important cause of acute lower respiratory tract disease and mortality for infants and children in tropical regions of the world.

From seroprevalence studies and studies of local epidemics, the initial *C. pneumoniae* infection usually happens between the ages of 5 and 20 years. In the USA, for example, there are

estimated to be 30,000 cases of childhood pneumonia each year caused by *C. pneumoniae*. Infections may cluster among groups of children or young adults (e.g., school pupils or military conscripts).

C. pneumoniae causes 10 to 25% of community-acquired lower respiratory tract infections (as reported from Sweden, Italy, Finland, and the USA). During an epidemic, *C. pneumoniae* infection may account for 50 to 60% of the cases of pneumonia. During these periods, also, more episodes of mixed infections with *S. pneumoniae* have been reported.

Reinfection during adulthood is common; the clinical presentation tends to be milder. Based on population seroprevalence studies, there tends to be increased exposure with age, which is particularly evident among men. Some investigators have speculated that a persistent, asymptomatic *C. pneumoniae* infection state is common.

In adults of middle age or older, *C. pneumoniae* infection may progress to chronic bronchitis and sinusitis. A study in the USA revealed that the incidence of pneumonia caused by *C. pneumoniae* in persons younger than 60 years is 1 case per 1,000 persons per year; but in the elderly, the disease incidence rose three-fold. *C. pneumoniae* infection rarely leads to hospitalization, except in patients with an underlying illness.

Of considerable importance is the association of atherosclerosis and *C. pneumoniae* infection. There are several epidemiological studies showing a correlation of previous infections with *C. pneumoniae* and heart attacks, coronary artery and carotid artery disease. See, Saikku *et al.*, *Lancet* 2: 983 (1988); Thom *et al.*, *JAMA* 268: 68 (1992); Linnanmaki *et al.*, *Circulation* 87: 1030 (1993); Saikku *et al.*, *Annals Int. Med.* 116: 273 (1992); Melnick *et al.*, *Am. J. Med.* 95: 499 (1993). Moreover, the organisms has been detected in atheromas and fatty streaks of the coronary, carotid, peripheral arteries and aorta. See, Shor *et al.*, *South African Med. J.* 82: 158 (1992); Kuo *et al.*, *J. Infect. Dis.* 167: 841 (1993); Kuo *et al.*, *Arteriosclerosis and Thrombosis* 13: 1500 (1993); Campbell *et al.*, *J. Infect. Dis.* 172: 585 (1995); Chiu *et al.*, *Circulation* 96: 2144-2148 (1997). Viable *C. pneumoniae* has been recovered from the coronary and carotid artery. Ramirez *et al.*, *Annals Int. Med.* 125: 979 (1996); Jackson *et al.*, Abst. K121, p272, 36th ICAAC, New Orleans (1996). Furthermore, it has been shown that *C. pneumoniae* can induce changes of atherosclerosis in a rabbit model. See, Fong *et al.*, (1997) *Journal of Clinical*

Microbiology 35: 48. Taken together, these results indicate that it is highly probable that *C. pneumoniae* can cause atherosclerosis in humans, though the epidemiological importance of chlamydial atherosclerosis remains to be demonstrated.

A number of recent studies have also indicated an association between *C. pneumoniae* infection and asthma. Infection has been linked to wheezing, asthmatic bronchitis, adult-onset asthma and acute exacerbation of asthma in adults, and small-scale studies have shown that prolonged antibiotic treatment was effective at greatly reducing the severity of the disease in some individuals. Hahn *et al.*, *Ann Allergy Asthma Immunol.* 80: 45-49 (1998); Hahn *et al.*, *Epidemiol Infect.* 117: 513-517 (1996); Bjornsson *et al.*, *Scand J Infect. Dis.* 28: 63-69 (1996); Hahn, *J. Fam. Pract.* 41: 345-351 (1995); Allegra *et al.*, *Eur. Respir. J.* 7: 2165-2168 (1994); Hahn *et al.*, *JAMA* 266: 225-230 (1991).

In light of these results, a protective vaccine against disease caused by *C. pneumoniae* infection would be of considerable importance. There is not yet an effective vaccine for human *C. pneumoniae* infection. Nevertheless, studies with *C. trachomatis* and *C. psittaci* indicate that this is an attainable goal. For example, mice which have recovered from a lung infection with *C. trachomatis* are protected from infertility induced by a subsequent vaginal challenge. Pal *et al.*, *Infection and Immunity* 64: 5341 (1996). Similarly, sheep immunized with inactivated *C. psittaci* were protected from subsequent chlamydial-induced abortions and stillbirths. Jones *et al.*, *Vaccine* 13: 715 (1995). Protection from chlamydial infections has been associated with Th1 immune responses, particularly the induction of INF γ -producing CD4⁺ T cells. Igietsemes *et al.*, *Immunology* 5: 317 (1993). The adoptive transfer of CD4⁺ cell lines or clones to nude or SCID mice conferred protection from challenge or cleared chronic disease (Igietseme *et al.*, *Regional Immunology* 5: 317 (1993); Magee *et al.*, *Regional Immunology* 5: 305 (1993)), and *in vivo* depletion of CD4⁺ T cells exacerbated disease post-challenge (Landers *et al.*, *Infection & Immunity* 59: 3774 (1991); Magee *et al.*, *Infection & Immunity* 63: 516 (1995)). However, the presence of sufficiently high titres of neutralizing antibody at mucosal surfaces can also exert a protective effect. Cotter *et al.*, *Infection and Immunity* 63: 4704 (1995).

The extent of antigenic variation within the species *C. pneumoniae* is not well characterized. Serovars of *C. trachomatis* are defined on the basis of antigenic variation in major

outer membrane proteins (MOMP), but published *C. pneumoniae* MOMP gene sequences show no variation between several diverse isolates of the organism. See, Campbell *et al.*, *Infection and Immunity* 58: 93 (1990); McCafferty *et al.*, *Infection and Immunity* 63: 2387-9 (1995); Knudsen *et al.*, Third Meeting of the European Society for Chlamydia Research, Vienna (1996). Regions of the protein known to be conserved in other chlamydial MOMPs are conserved in *C. pneumoniae*. See, Campbell *et al.*, *Infection and Immunity* 58: 93 (1990); McCafferty *et al.*, *Infection and Immunity* 63: 2387-9 (1995). One study has described a strain of *C. pneumoniae* with a MOMP of greater than usual molecular weight, but the gene for this has not been sequenced. Grayston *et al.*, *J. Infect. Dis.* 168: 1231 (1995). Partial sequences of outer membrane protein 2 from nine diverse isolates were also found to be invariant. Ramirez *et al.*, *Annals Int. Med.* 125: 979 (1996). The genes for HSP60 and HSP70 show little variation from other chlamydial species, as would be expected. The gene encoding a 76 kDa antigen has been cloned from a single strain of *C. pneumoniae*. It has no significant similarity with other known chlamydial genes. Marrie, *Clin. Infect. Dis.* 18: 501 (1993).

Many antigens recognized by immune sera to *C. pneumoniae* are conserved across all chlamydiae, but 98kDa, 76 kDa and 54 kDa proteins may be *C. pneumoniae*-specific. Campos *et al.*, *Invest. Ophthalmol. Vis. Sci.* 36: 1477 (1995); Marrie, *Clin. Infect. Dis.* 18: 501 (1993); Wiedmann-Al-Ahmad *et al.*, *Clin. Diagn. Lab. Immunol.* 4: 700-704 (1997). Immunoblotting of isolates with sera from patients does show variation of blotting patterns between isolates, indicating that serotypes *C. pneumoniae* may exist. Grayston *et al.*, *J. Infect. Dis.* 168: 1231 (1995); Ramirez *et al.*, *Annals Int. Med.* 125: 979 (1996). However, the results are potentially confounded by the infection status of the patients, since immunoblot profiles of a patient's sera change with time post-infection. An assessment of the number and relative frequency of any serotypes, and the defining antigens, is not yet possible.

Thus, a need remains for effective compositions for preventing, treating, and diagnosing *Chlamydia* infections.

SUMMARY OF THE INVENTION

In one aspect, the present invention provides purified and isolated DNA molecules that encode *Chlamydia* which can be used in methods to prevent, treat, and diagnose *Chlamydia* infection. Encoded polypeptides, designated 98 kDa putative outer membrane protein, include polypeptides having the amino acid sequence shown in SEQ ID NO: 2 and the DNA molecules include SEQ ID NO: 1 full-length sequence (top sequence) and coding sequence (bottom sequence) for the mature polypeptide. Those skilled in the art will appreciate that the invention also includes DNA molecules that encode mutants, variants, and derivatives of such polypeptides, which result from the addition, deletion, or substitution of non-essential amino acids as described herein. The invention also includes RNA molecules corresponding to the DNA molecules of the invention.

In addition to the DNA and RNA molecules, the invention includes the corresponding polypeptides and monospecific antibodies that specifically bind to such polypeptides.

The present invention has wide application and includes expression cassettes, vectors, and cells transformed or transfected with the polynucleotides of the invention. Accordingly, the present invention provides (i) a method for producing a polypeptide of the invention in a recombinant host system and related expression cassettes, vectors, and transformed or transfected cells; (ii) a live vaccine vectors such as viral or bacterial live vaccine vectors, including, pox virus, alphavirus, *Salmonella typhimurium*, or *Vibrio cholerae* vector, containing a polynucleotide of the invention, such vaccine vectors being useful for, e.g., preventing and treating *Chlamydia* infection, in combination with a diluent or carrier, and related pharmaceutical compositions and associated therapeutic and/or prophylactic methods; (iii) a therapeutic and/or prophylactic method involving administration of an RNA or DNA molecule of the invention, either in a naked form or formulated with a delivery vehicle, a polypeptide or combination of polypeptides, or a monospecific antibody of the invention, and related pharmaceutical compositions; (iv) a method for diagnosing the presence of *Chlamydia* in a biological sample, which can involve the use of a DNA or RNA molecule, a monospecific antibody, or a polypeptide of the invention; and (v) a method for purifying a polypeptide of the invention by antibody-based affinity chromatography. The present invention provides purified and isolated

DNA molecules, which encode *Chlamydia* that can be used in methods to prevent, treat, and diagnose *Chlamydia* infection.

BRIEF DESCRIPTION OF THE DRAWINGS

The present invention will be further understood from the following description with
5 reference to the drawings, in which:

FIG. 1 shows the nucleotide sequence (top sequence) and the deduced amino acid sequence (bottom sequence) of the full length 98 kDa putative outer membrane protein gene (SEQ ID NO: 1) and the processed sequence from *Chlamydia pneumoniae* (SEQ ID NO: 2).

FIG. 2 shows the restriction enzyme analysis of nucleotide sequence encoding the
10 *C. pneumoniae* 98 kDa putative outer membrane protein gene.

FIG. 3 shows the construction and elements of plasmid pCAI396.

FIG. 4 illustrates protection against *C. pneumoniae* infection by pCAI396 following DNA immunization. Individual data points are shown for each animal (hollow diamonds) as well as mean and standard deviation for each group (solid squares).

DETAILED DESCRIPTION OF THE INVENTION

In the *C. pneumoniae* genome, open reading frames (ORFs) encoding chlamydial polypeptides have been identified. These polypeptides include polypeptides permanently found in the bacterial membrane structure, polypeptides that are present in the external vicinity of the bacterial membrane, include polypeptides permanently found in the inclusion membrane
20 structure, polypeptides that are present in the external vicinity of the inclusion membrane, and polypeptides that are released into the cytoplasm of the infected cell. These polypeptides can be used in vaccination methods for preventing and treating *Chlamydia* infection.

According to a first aspect of the invention, there are provided isolated polynucleotides encoding the precursor and mature forms of *Chlamydia* polypeptides.

An isolated polynucleotide of the invention encodes a polypeptide having an amino acid sequence that is homologous to a *Chlamydia* amino acid sequence, the *Chlamydia* amino acid sequence being selected from the group consisting of the amino acid sequences as shown in SEQ ID NOS: 1 and 2.

5 The term "isolated polynucleotide" is defined as a polynucleotide removed from the environment in which it naturally occurs. For example, a naturally-occurring DNA molecule present in the genome of a living bacteria or as part of a gene bank is not isolated, but the same molecule separated from the remaining part of the bacterial genome, as a result of, *e.g.*, a cloning event (amplification), is isolated. Typically, an isolated DNA molecule is free from DNA
10 regions (*e.g.*, coding regions) with which it is immediately contiguous at the 5' or 3' end, in the naturally occurring genome. Such isolated polynucleotides could be part of a vector or a composition and still be isolated in that such a vector or composition is not part of its natural environment.

A polynucleotide of the invention can be in the form of RNA or DNA (*e.g.*, cDNA,
15 genomic DNA, or synthetic DNA), or modifications or combinations thereof. The DNA can be double-stranded or single-stranded, and, if single-stranded, can be the coding strand or the non-coding (anti-sense) strand. The sequence that encodes a polypeptide of the invention as shown in SEQ ID NO: 1 can be (a) the coding sequence (bottom sequence); (b) a ribonucleotide sequence derived by transcription of (a); or (c) a different coding sequence. This latter, as a
20 result of the redundancy or degeneracy of the genetic code, encodes the same polypeptides as the DNA molecules of which the nucleotide sequences are illustrated in SEQ ID NOS: 1 or 2.

By "homologous amino acid sequence" is meant an amino acid sequence that differs from an amino acid sequence shown in SEQ ID NO: 2, only by one or more conservative amino acid substitutions, or by one or more non-conservative amino acid substitutions, deletions, or
25 additions located at positions at which they do not destroy the specific antigenicity of the polypeptide.

Preferably, such a sequence is at least 75%, more preferably 80%, and most preferably 90% identical to an amino acid sequence shown in SEQ ID NO: 2.

Homologous amino acid sequences include sequences that are identical or substantially identical to an amino acid sequence as shown in SEQ ID NO: 2. By "amino acid sequence substantially identical" is meant a sequence that is at least 90%, preferably 95%, more preferably 97%, and most preferably 99% identical to an amino acid sequence of reference and that
5 preferably differs from the sequence of reference, if at all, by a majority of conservative amino acid substitutions.

Conservative amino acid substitutions typically include substitutions among amino acids of the same class. These classes include, for example, (a) amino acids having uncharged polar side chains, such as asparagine, glutamine, serine, threonine, and tyrosine; (b) amino acids
10 having basic side chains, such as lysine, arginine, and histidine; (c) amino acids having acidic side chains, such as aspartic acid and glutamic acid; and (d) amino acids having nonpolar side chains, such as glycine, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan, and cysteine.

Homology is typically measured using sequence analysis software (*e.g.*, Sequence
15 Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705). Similar amino acid sequences are aligned to obtain the maximum degree of homology (*i.e.*, identity). To this end, it may be necessary to artificially introduce gaps into the sequence. Once the optimal alignment has been set up, the degree of homology (*i.e.*, identity) is established by recording all of the
20 positions in which the amino acids of both sequences are identical, relative to the total number of positions.

Similarity factors include similar size, shape and electrical charge. One particularly preferred method of determining amino acid similarities is the PAM250 matrix described in Dayhoff et al., 5 ATLAS OF PROTEIN SEQUENCE AND STRUCTURE 345-352 (1978 & Supp.),
25 incorporated by reference herein. A similarity score is first calculated as the sum of the aligned pairwise amino acid similarity scores. Insertions and deletions are ignored for the purposes of percent homology and identity. Accordingly, gap penalties are not used in this calculation. The raw score is then normalized by dividing it by the geometric mean of the scores of the candidate

compound and the reference sequence. The geometric mean is the square root of the product of these scores. The normalized raw score is the percent homology.

Preferably, a homologous sequence is one that is at least 45%, more preferably 60%, and most preferably 85% identical to the coding sequence of SEQ ID NO: 1.

5 Polypeptides having a sequence homologous to one of the sequences shown in SEQ ID NOS: 1 and 2, include naturally-occurring allelic variants, as well as mutants and variants or any other non-naturally-occurring variants that are analogous in terms of antigenicity, to a polypeptide having a sequence as shown in SEQ ID NOS: 1 or 2.

10 An allelic variant is an alternate form of a polypeptide that is characterized as having a substitution, deletion, or addition of one or more amino acids that does not substantially alter the biological function of the polypeptide. By "biological function" is meant the function of the polypeptide in the cells in which it naturally occurs, even if the function is not necessary for the growth or survival of the cells. For example, the biological function of a porin is to allow the entry into cells of compounds present in the extracellular medium. The biological function is
15 distinct from the antigenic function. A polypeptide can have more than one biological function.

Allelic variants are very common in nature. For example, a bacterial species, *e.g.*, *C. pneumoniae*, is usually represented by a variety of strains that differ from each other by minor allelic variations. Indeed, a polypeptide that fulfills the same biological function in different strains can have an amino acid sequence that is not identical in each of the strains. Such an
20 allelic variation may be equally reflected at the polynucleotide level.

Support for the use of allelic variants of polypeptide antigens comes from, *e.g.*, studies of the *Chlamydial* MOMP antigen. The amino acid sequence of the MOMP varies from strain to strain, yet cross-strain antibody binding plus neutralization of infectivity occurs, indicating that the MOMP, when used as an immunogen, is tolerant of amino acid variations.

25 Polynucleotides, *e.g.*, DNA molecules, encoding allelic variants can easily be retrieved by polymerase chain reaction (PCR) amplification of genomic bacterial DNA extracted by conventional methods. This involves the use of synthetic oligonucleotide primers matching upstream and downstream of the 5' and 3' ends of the encoding domain. Suitable primers can be

designed according to the nucleotide sequence information provided in SEQ ID NOS: 1 and 2. Typically, a primer can consist of 10 to 40, preferably 15 to 25 nucleotides. It may be also advantageous to select primers containing C and G nucleotides in a proportion sufficient to ensure efficient hybridization; *e.g.*, an amount of C and G nucleotides of at least 40%, preferably
5 50% of the total nucleotide amount.

Useful homologs that do not naturally occur can be designed using known methods for identifying regions of an antigen that are likely to be tolerant of amino acid sequence changes and/or deletions. For example, sequences of the antigen from different species can be compared to identify conserved sequences.

10 Polypeptide derivatives that are encoded by polynucleotides of the invention include, *e.g.*, fragments, polypeptides having large internal deletions derived from full-length polypeptides, and fusion proteins.

Polypeptide fragments of the invention can be derived from a polypeptide having a sequence homologous to any of the sequences shown in SEQ ID NOS: 1 and 2, to the extent that
15 the fragments retain the desired substantial antigenicity of the parent polypeptide (specific antigenicity). Polypeptide derivatives can also be constructed by large internal deletions that remove a substantial part of the parent polypeptide, while retaining the desired specific antigenicity. Generally, polypeptide derivatives should be about at least 12 amino acids in length to maintain the antigenicity. Advantageously, they can be at least 20 amino acids, preferably at
20 least 50 amino acids, more preferably at least 75 amino acids, and most preferably at least 100 amino acids in length.

Useful polypeptide derivatives, *e.g.*, polypeptide fragments, can be designed using computer-assisted analysis of amino acid sequences in order to identify sites in protein antigens having potential as surface-exposed, antigenic regions. Hughes *et al.*, *Infect. Immun.* 60: 3497
25 (1992).

Polypeptide fragments and polypeptides having large internal deletions can be used for revealing epitopes that are otherwise masked in the parent polypeptide and that may be of

importance for inducing, for example, a protective T cell-dependent immune response. Deletions can also remove immunodominant regions of high variability among strains.

It is an accepted practice in the field of immunology to use fragments and variants of protein immunogens as vaccines and immunogens, as all that is required to induce an immune response to a protein may be a small (*e.g.*, 8 to 10 amino acid) region of the protein. This has been done for a number of vaccines against pathogens other than *Chlamydia*. For example, short synthetic peptides corresponding to surface-exposed antigens of pathogens such as murine mammary tumor virus, peptide containing 11 amino acids (Dion *et al.*, *Virology* 179: 474-477 (1990)); Semliki Forest virus, peptide containing 16 amino acids (Snijders *et al.*, *J. Gen. Virol.* 72: 557-565 (1991)); and canine parvovirus, two overlapping peptides, each containing 15 amino acids (Langeveld *et al.*, *Vaccine* 12: 1473-1480 (1994)) have been shown to be effective vaccine antigens against their respective pathogens.

Polynucleotides encoding polypeptide fragments and polypeptides having large internal deletions can be constructed using standard methods (see, *e.g.*, Ausubel *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons Inc. (1994)); for example, by PCR, including inverse PCR, by restriction enzyme treatment of the cloned DNA molecules, or by the method of Kunkel *et al.* (*Proc. Natl. Acad. Sci. USA* 82: 448 (1985)); biological material available at Stratagene.

A polypeptide derivative can also be produced as a fusion polypeptide that contains a polypeptide or a polypeptide derivative of the invention fused, *e.g.*, at the N- or C-terminal end, to any other polypeptide. For construction of DNA encoding the amino acid sequence corresponding to hybrid fusion proteins, a first DNA encoding amino acid sequence corresponding to portions of SEQ ID NO: 1 or 2 is joined to a second DNA using methods described in, for example, U.S. Patent 5,844,095, incorporated herein by reference. A product can then be easily obtained by translation of the genetic fusion. Vectors for expressing fusion polypeptides are commercially available, such as the pMal-c2 or pMal-p2 systems of New England Biolabs, in which the fusion peptide is a maltose binding protein, the glutathione-S-transferase system of Pharmacia, or the His-Tag system available from Novagen.

These and other expression systems provide convenient means for further purification of polypeptides and derivatives of the invention.

Another particular example of fusion polypeptides included in the invention includes a polypeptide or polypeptide derivative of the invention fused to a polypeptide having adjuvant activity, such as, *e.g.*, the subunit B of either cholera toxin or *E. coli* heat-labile toxin. Several possibilities are can be used for achieving fusion. First, the polypeptide of the invention can be fused to the N-, or preferably, to the C-terminal end of the polypeptide having adjuvant activity. Second, a polypeptide fragment of the invention can be fused within the amino acid sequence of the polypeptide having adjuvant activity.

As stated above, the polynucleotides of the invention encode *Chlamydia* polypeptides in precursor or mature form. They can also encode hybrid precursors containing heterologous signal peptides, which can mature into polypeptides of the invention. By "heterologous signal peptide" is meant a signal peptide that is not found in the naturally-occurring precursor of a polypeptide of the invention.

A polynucleotide of the invention, having a homologous coding sequence, hybridizes, preferably under stringent conditions, to a polynucleotide having a sequence as shown in SEQ ID NO: 1. Hybridization procedures are described in, *e.g.*, Ausubel *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons Inc. (1994); Silhavy *et al.*, EXPERIMENTS WITH GENE FUSIONS, Cold Spring Harbor Laboratory Press (1984); Davis *et al.*, A MANUAL FOR GENETIC ENGINEERING: ADVANCED BACTERIAL GENETICS, Cold Spring Harbor Laboratory Press (1980), each incorporated herein by reference. Important parameters that can be considered for optimizing hybridization conditions are reflected in a formula that allows calculation of a critical value, the melting temperature above which two complementary DNA strands separate from each other. Casey and Davidson, *Nucl. Acid Res.* 4: 1539 (1977). This formula is as follows:

$$T_m = 81.5 + 0.5 \times (\% \text{ G+C}) + 1.6 \log (\text{positive ion concentration}) - 0.6 \times (\% \text{ formamide}).$$

Under appropriate stringency conditions, hybridization temperature (T_h) is approximately 20-40°C, 20-25°C or, preferably, 30-40°C below the calculated T_m . Those skilled in the art will

understand that optimal temperature and salt conditions can be readily determined empirically in preliminary experiments using conventional procedures.

For example, stringent conditions can be achieved, both for pre-hybridizing and hybridizing incubations, (i) within 4-16 hours at 42°C, in 6xSSC containing 50% formamide or
5 (ii) within 4-16 hours at 65°C in an aqueous 6xSSC solution (1 M NaCl, 0.1 M sodium citrate (pH 7.0)).

For polynucleotides containing 30 to 600 nucleotides, the above formula is used and then is corrected by subtracting (600/polynucleotide size in base pairs). Stringency conditions are defined by a T_h that is 5 to 10°C below T_m .

10 Hybridization conditions with oligonucleotides shorter than 20-30 bases do not exactly follow the rules set forth above. In such cases, the formula for calculating the T_m is as follows:

$$T_m = 4 \times (G+C) + 2 (A+T).$$

For example, an 18 nucleotide fragment of 50% G+C would have an approximate T_m of 54°C.

A polynucleotide molecule of the invention, containing RNA, DNA, or modifications or
15 combinations thereof, can have various applications. For example, a DNA molecule can be used (i) in a process for producing the encoded polypeptide in a recombinant host system, (ii) in the construction of vaccine vectors such as poxviruses, which are further used in methods and compositions for preventing and/or treating *Chlamydia* infection, (iii) as a vaccine agent (as well as an RNA molecule), in a naked form or formulated with a delivery vehicle and, (iv) in the
20 construction of attenuated *Chlamydia* strains that can overexpress a polynucleotide of the invention or express it in a modified, mutated form, such as a non-toxic form, if appropriate.

For vaccine compositions and uses of the proteins and peptides and encoding nucleotides of the present invention for protection against diseases caused by *Chlamydia*, it is not preferred to use naked DNA encoding the protein or peptides and administering these nucleotides intranasally or
25 intramuscularly. For these proteins, it is preferred to administer the encoding nucleic acids by other routes such as intradermally and/or to formulate the encoding nucleic acids to improve (or adjuvant) the immune response. It is also preferred to include the encoding nucleic acid as part of a recombinant live vector, such as a viral or bacterial vector for use as the immunizing agent.

It is also preferred to immunize with vaccine formulations comprising the proteins or peptides of the invention themselves. These vaccine formulations may include the use of adjuvants.

According to a second aspect of the invention, there is therefore provided (i) an expression cassette containing a DNA molecule of the invention placed under the control of the elements required for expression, in particular under the control of an appropriate promoter;
5 (ii) an expression vector containing an expression cassette of the invention; (iii) a prokaryotic or eukaryotic cell transformed or transfected with an expression cassette and/or vector of the invention, as well as (iv) a process for producing a polypeptide or polypeptide derivative encoded by a polynucleotide of the invention, which involves culturing a prokaryotic or
10 eukaryotic cell transformed or transfected with an expression cassette and/or vector of the invention, under conditions that allow expression of the DNA molecule of the invention and, recovering the encoded polypeptide or polypeptide derivative from the cell culture.

A recombinant expression system can be selected from prokaryotic and eukaryotic hosts. Eukaryotic hosts include yeast cells (*e.g.*, *Saccharomyces cerevisiae* or *Pichia pastoris*),
15 mammalian cells (*e.g.*, COS1, NIH3T3, or JEG3 cells), arthropods cells (*e.g.*, *Spodoptera frugiperda* (SF9) cells), and plant cells. Preferably, a prokaryotic host such as *E. coli* is used. Bacterial and eukaryotic cells are available from a number of different sources to those skilled in the art, *e.g.*, the American Type Culture Collection (ATCC; Rockville, Maryland).

The choice of the expression system depends on the features desired for the expressed
20 polypeptide. For example, it may be useful to produce a polypeptide of the invention in a particular lipidated form or any other form.

The choice of the expression cassette will depend on the host system selected as well as the features desired for the expressed polypeptide. Typically, an expression cassette includes a promoter that is functional in the selected host system and can be constitutive or inducible; a
25 ribosome binding site; a start codon (ATG) if necessary, a region encoding a signal peptide, *e.g.*, a lipidation signal peptide; a DNA molecule of the invention; a stop codon; and optionally a 3' terminal region (translation and/or transcription terminator). The signal peptide encoding region is adjacent to the polynucleotide of the invention and placed in proper reading frame. The signal peptide-encoding region can be homologous or heterologous to the DNA molecule encoding the

mature polypeptide and can be specific to the secretion apparatus of the host used for expression.

The open reading frame constituted by the DNA molecule of the invention, solely or together with the signal peptide, is placed under the control of the promoter so that transcription and translation occur in the host system. Promoters, signal peptide encoding regions are widely

5 known and available to those skilled in the art and includes, for example, the promoter of *Salmonella typhimurium* (and derivatives) that is inducible by arabinose (promoter araB) and is functional in Gram-negative bacteria such as *E. coli* (as described in U.S. Patent 5,028,530 and in Cagnon *et al.*, *Protein Engineering* 4: 843 (1991)); the promoter of the gene of bacteriophage T7 encoding RNA polymerase, that is functional in a number of *E. coli* strains expressing T7
10 polymerase (described in U.S. Patent 4,952,496); OspA lipidation signal peptide; and RlpB lipidation signal peptide (Takase *et al.*, *J. Bact.* 169: 5692 (1987)).

The expression cassette is typically part of an expression vector, which is selected for its ability to replicate in the chosen expression system. Expression vectors (*e.g.*, plasmids or viral vectors) can be chosen from those described in Pouwels *et al.* (CLONING VECTORS:

15 LABORATORY MANUAL, 85, Supp. 1987). They can be purchased from various commercial sources.

Methods for transforming/transfecting host cells with expression vectors will depend on the host system selected as described in Ausubel *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons Inc. (1994).

20 Upon expression, a recombinant polypeptide of the invention (or a polypeptide derivative) is produced and remains in the intracellular compartment, is secreted/excreted in the extracellular medium or in the periplasmic space, or is embedded in the cellular membrane. The polypeptide can then be recovered in a substantially purified form from the cell extract or from the supernatant after centrifugation of the recombinant cell culture. Typically, the recombinant
25 polypeptide can be purified by antibody-based affinity purification or by any other method that can be readily adapted by a person skilled in the art, such as by genetic fusion to a small affinity binding domain. Antibody-based affinity purification methods are also available for purifying a polypeptide of the invention extracted from a *Chlamydia* strain. Antibodies useful for purifying by immunoaffinity the polypeptides of the invention can be obtained as described below.

A polynucleotide of the invention can also be useful in the vaccine field, *e.g.*, for achieving DNA vaccination. There are two major possibilities, either using a viral or bacterial host as gene delivery vehicle (live vaccine vector) or administering the gene in a free form, *e.g.*, inserted into a plasmid. Therapeutic or prophylactic efficacy of a polynucleotide of the invention
5 can be evaluated as described below.

Accordingly, in a third aspect of the invention, there is provided (i) a vaccine vector such as a poxvirus, containing a DNA molecule of the invention, placed under the control of elements required for expression; (ii) a composition of matter containing a vaccine vector of the invention, together with a diluent or carrier; particularly, (iii) a pharmaceutical composition containing a
10 therapeutically or prophylactically effective amount of a vaccine vector of the invention; (iv) a method for inducing an immune response against *Chlamydia* in a mammal (*e.g.*, a human; alternatively, the method can be used in veterinary applications for treating or preventing *Chlamydia* infection of animals, *e.g.*, cats or birds), which involves administering to the mammal an immunogenically effective amount of a vaccine vector of the invention to elicit an immune
15 response, *e.g.*, a protective or therapeutic immune response to *Chlamydia*; and particularly, (v) a method for preventing and/or treating a *Chlamydia* (*e.g.*, *C. trachomatis*, *C. psittaci*, *C. pneumonia*, *C. pecorum*) infection, which involves administering a prophylactic or therapeutic amount of a vaccine vector of the invention to an individual in need. Additionally, the third aspect of the invention encompasses the use of a vaccine vector of the invention in the
20 preparation of a medicament for preventing and/or treating *Chlamydia* infection.

A vaccine vector of the invention can express one or several polypeptides or derivatives of the invention, as well as at least one additional *Chlamydia* antigen, fragment, homolog, mutant, or derivative thereof. In addition, it can express a cytokine, such as interleukin-2 (IL-2) or interleukin-12 (IL-12), that enhances the immune response (adjuvant effect). Thus, a vaccine
25 vector can include an additional DNA sequence encoding, *e.g.*, a chlamydial antigen, or a cytokine, placed under the control of elements required for expression in a mammalian cell.

Alternatively, a composition of the invention can include several vaccine vectors, each of them being capable of expressing a polypeptide or derivative of the invention. A composition

can also contain a vaccine vector capable of expressing an additional *Chlamydia* antigen, or a subunit, fragment, homolog, mutant, or derivative thereof; or a cytokine such as IL-2 or IL-12.

In vaccination methods for treating or preventing infection in a mammal, a vaccine vector of the invention can be administered by any conventional route in use in the vaccine field, particularly, to a mucosal (*e.g.*, ocular, intranasal, oral, gastric, pulmonary, intestinal, rectal, vaginal, or urinary tract) surface or *via* the parenteral (*e.g.*, subcutaneous, intradermal, intramuscular, intravenous, or intraperitoneal) route. Preferred routes depend upon the choice of the vaccine vector. The administration can be achieved in a single dose or repeated at intervals. The appropriate dosage depends on various parameters understood by skilled artisans such as the vaccine vector itself, the route of administration or the condition of the mammal to be vaccinated (weight, age and the like).

Live vaccine vectors available in the art include viral vectors such as adenoviruses, alphavirus, and poxviruses as well as bacterial vectors, *e.g.*, *Shigella*, *Salmonella*, *Vibrio cholerae*, *Lactobacillus*, Bacille bilié de Calmette-Guérin (BCG), and *Streptococcus*.

An example of an adenovirus vector, as well as a method for constructing an adenovirus vector capable of expressing a DNA molecule of the invention, are described in U.S. Patent 4,920,209. Poxvirus vectors that can be used include, *e.g.*, vaccinia and canary pox virus, described in U.S. Patent 4,722,848 and U.S. Patent 5,364,773, respectively (also see, *e.g.*, Tartaglia *et al.*, *Virology* 188: 217 (1992)) for a description of a vaccinia virus vector; and Taylor *et al.*, *Vaccine* 13: 539 (1995) for a reference of a canary pox). Poxvirus vectors capable of expressing a polynucleotide of the invention can be obtained by homologous recombination as described in Kieny *et al.*, *Nature* 312: 163 (1984) so that the polynucleotide of the invention is inserted in the viral genome under appropriate conditions for expression in mammalian cells. Generally, the dose of vaccine viral vector, for therapeutic or prophylactic use, can be of from about 1×10^4 to about 1×10^{11} , advantageously from about 1×10^7 to about 1×10^{10} , preferably of from about 1×10^7 to about 1×10^9 plaque-forming units per kilogram. Preferably, viral vectors are administered parenterally; for example, in three doses, four weeks apart. Those skilled in the art recognize that it is preferable to avoid adding a chemical adjuvant to a composition containing a

viral vector of the invention and thereby minimizing the immune response to the viral vector itself.

Non-toxicogenic *Vibrio cholerae* mutant strains that are useful as a live oral vaccine are described in Mekalanos *et al.*, *Nature* 306: 551 (1983) and U.S. Patent 4,882,278 (strain in which a substantial amount of the coding sequence of each of the two *ctxA* alleles has been deleted so that no functional *cholerae* toxin is produced); WO 92/11354 (strain in which the *irgA* locus is inactivated by mutation; this mutation can be combined in a single strain with *ctxA* mutations); and WO 94/1533 (deletion mutant lacking functional *ctxA* and *attRS1* DNA sequences). These strains can be genetically engineered to express heterologous antigens, as described in WO 94/19482. An effective vaccine dose of a *Vibrio cholerae* strain capable of expressing a polypeptide or polypeptide derivative encoded by a DNA molecule of the invention can contain, e.g., about 1×10^5 to about 1×10^9 , preferably about 1×10^6 to about 1×10^8 viable bacteria in an appropriate volume for the selected route of administration. Preferred routes of administration include all mucosal routes; most preferably, these vectors are administered intranasally or orally.

Attenuated *Salmonella typhimurium* strains, genetically engineered for recombinant expression of heterologous antigens or not, and their use as oral vaccines are described in Nakayama *et al.*, *Bio/Technology* 6: 693 (1988) and WO 92/11361. Preferred routes of administration include all mucosal routes; most preferably, these vectors are administered intranasally or orally.

Others bacterial strains useful as vaccine vectors are described in High *et al.*, *EMBO* 11: 1991 (1992); Sizemore *et al.*, *Science* 270: 299 (1995) (*Shigella flexneri*); Medaglini *et al.*, *Proc. Natl. Acad. Sci. USA* 92: 6868 (1995) (*Streptococcus gordonii*); and Flynn, *Cell. Mol. Biol.* 40: 31 (1994), WO 88/6626, WO 90/0594, WO 91/13157, WO 92/1796, and WO 92/21376 (Bacille Calmette Guerin).

In bacterial vectors, polynucleotide of the invention can be inserted into the bacterial genome or can remain in a free state, carried on a plasmid.

An adjuvant can also be added to a composition containing a vaccine bacterial vector. A number of adjuvants are known to those skilled in the art. Preferred adjuvants can be selected from the list provided below.

According to a fourth aspect of the invention, there is also provided (i) a composition of matter containing a polynucleotide of the invention, together with a diluent or carrier; (ii) a pharmaceutical composition containing a therapeutically or prophylactically effective amount of a polynucleotide of the invention; (iii) a method for inducing an immune response against *Chlamydia*, in a mammal, by administering to the mammal, an immunogenically effective amount of a polynucleotide of the invention to elicit an immune response, *e.g.*, a protective immune response to *Chlamydia*; and particularly, (iv) a method for preventing and/or treating a *Chlamydia* (*e.g.*, *C. trachomatis*, *C. psittaci*, *C. pneumoniae*, or *C. pecorum*) infection, by administering a prophylactic or therapeutic amount of a polynucleotide of the invention to an individual in need. Additionally, the fourth aspect of the invention encompasses the use of a polynucleotide of the invention in the preparation of a medicament for preventing and/or treating *Chlamydia* infection. The fourth aspect of the invention preferably includes the use of a DNA molecule placed under conditions for expression in a mammalian cell, *e.g.*, in a plasmid that is unable to replicate in mammalian cells and to substantially integrate in a mammalian genome.

Polynucleotides (DNA or RNA) of the invention can also be administered as such to a mammal for vaccine, *e.g.*, therapeutic or prophylactic, purpose. When a DNA molecule of the invention is used, it can be in the form of a plasmid that is unable to replicate in a mammalian cell and unable to integrate in the mammalian genome. Typically, a DNA molecule is placed under the control of a promoter suitable for expression in a mammalian cell. The promoter can function ubiquitously or tissue-specifically. Examples of non-tissue specific promoters include the early Cytomegalovirus (CMV) promoter (described in U.S. Patent 4,168,062) and the Rous Sarcoma Virus promoter (described in Norton & Coffin, *Molec. Cell Biol.* 5: 281(1985)). The desmin promoter (Li *et al.*, *Gene* 78: 243 (1989), Li & Paulin, *J. Biol. Chem.* 266: 6562 (1991), and Li & Paulin, *J. Biol. Chem.* 268: 10403 (1993)) is tissue-specific and drives expression in muscle cells. More generally, useful vectors are described, *i.a.*, WO 94/21797 and Hartikka *et al.*, *Human Gene Therapy* 7: 1205 (1996).

For DNA/RNA vaccination, the polynucleotide of the invention can encode a precursor or a mature form. When it encodes a precursor form, the precursor form can be homologous or heterologous. In the latter case, a eukaryotic leader sequence can be used, such as the leader sequence of the tissue-type plasminogen factor (tPA).

5 A composition of the invention can contain one or several polynucleotides of the invention. It can also contain at least one additional polynucleotide encoding another *Chlamydia* antigen or a fragment, derivative, mutant, or analog thereof. A polynucleotide encoding a cytokine, such as interleukin-2 (IL-2) or interleukin-12 (IL-12), can also be added to the composition so that the immune response is enhanced. These additional polynucleotides are
10 placed under appropriate control for expression. Advantageously, DNA molecules of the invention and/or additional DNA molecules to be included in the same composition, can be carried in the same plasmid.

Standard techniques of molecular biology for preparing and purifying polynucleotides can be used in the preparation of polynucleotide therapeutics of the invention. For use as a
15 vaccine, a polynucleotide of the invention can be formulated according to various methods.

First, a polynucleotide can be used in a naked form, free of any delivery vehicles, such as anionic liposomes, cationic lipids, microparticles, *e.g.*, gold microparticles, precipitating agents, *e.g.*, calcium phosphate, or any other transfection-facilitating agent. In this case, the polynucleotide can be simply diluted in a physiologically acceptable solution, such as sterile
20 saline or sterile buffered saline, with or without a carrier. When present, the carrier preferably is isotonic, hypotonic, or weakly hypertonic, and has a relatively low ionic strength, such as provided by a sucrose solution, *e.g.*, a solution containing 20% sucrose.

Alternatively, a polynucleotide can be associated with agents that assist in cellular uptake. It can be, *i.a.*, (i) complemented with a chemical agent that modifies the cellular permeability,
25 such as bupivacaine (see, *e.g.*, WO 94/16737), (ii) encapsulated into liposomes, or (iii) associated with cationic lipids or silica, gold, or tungsten microparticles.

Anionic and neutral liposomes are well-known in the art (see, *e.g.*, LIPOSOMES: A PRACTICAL APPROACH, RPC New Ed, IRL Press (1990)), for a detailed description of methods

for making liposomes) and are useful for delivering a large range of products, including polynucleotides.

Cationic lipids are also known in the art and are commonly used for gene delivery. Such lipids include Lipofectin™ also known as DOTMA

5 (N-[1-(2,3-dioleyloxy)propyl]-N,N,N-trimethylammonium chloride), DOTAP (1,2-bis(oleyloxy)-3-(trimethylammonio)propane), DDAB (dimethyldioctadecylammonium bromide), DOGS (dioctadecylamidologlycyl spermine) and cholesterol derivatives such as DC-Chol (3 beta-(N-(N',N'-dimethyl aminomethane)-carbamoyl) cholesterol). A description of these cationic lipids can be found in EP 187,702, WO 90/11092, U.S. Patent 5,283,185,
10 WO 91/15501, WO 95/26356, and U.S. Patent 5,527,928. Cationic lipids for gene delivery are preferably used in association with a neutral lipid such as DOPE (dioleoyl phosphatidylethanolamine), as described in, *e.g.*, WO 90/11092.

Other transfection-facilitating compounds can be added to a formulation containing cationic liposomes. A number of them are described in, *e.g.*, WO 93/18759, WO 93/19768,
15 WO 94/25608, and WO 95/2397. They include, *i.a.*, spermine derivatives useful for facilitating the transport of DNA through the nuclear membrane (see, for example, WO 93/18759) and membrane-permeabilizing compounds such as GALA, Gramicidine S, and cationic bile salts (see, for example, WO 93/19768).

Gold or tungsten microparticles can also be used for gene delivery, as described in
20 WO 91/359, WO 93/17706, and Tang *et al.* (*Nature* 356: 152 (1992)). In this case, the microparticle-coated polynucleotides can be injected *via* intradermal or intra-epidermal routes using a needleless injection device ("gene gun"), such as those described in U.S. Patent 4,945,050, U.S. Patent 5,015,580, and WO 94/24263.

The amount of DNA to be used in a vaccine recipient depends, *e.g.*, on the strength of the
25 promoter used in the DNA construct, the immunogenicity of the expressed gene product, the condition of the mammal intended for administration (*e.g.*, the weight, age, and general health of the mammal), the mode of administration, and the type of formulation. In general, a therapeutically or prophylactically effective dose from about 1 µg to about 1 mg, preferably, from about 10 µg to about 800 µg and, more preferably, from about 25 µg to about 250 µg, can

be administered to human adults. The administration can be achieved in a single dose or repeated at intervals.

The route of administration can be any conventional route used in the vaccine field. As general guidance, a polynucleotide of the invention can be administered *via* a mucosal surface, *e.g.*, an ocular, intranasal, pulmonary, oral, intestinal, rectal, vaginal, and urinary tract surface; or *via* a parenteral route, *e.g.*, by an intravenous, subcutaneous, intraperitoneal, intradermal, intra-epidermal, or intramuscular route. The choice of the administration route will depend on, *e.g.*, the formulation that is selected. A polynucleotide formulated in association with bupivacaine is advantageously administered into muscles. When a neutral or anionic liposome or a cationic lipid, such as DOTMA or DC-Chol, is used, the formulation can be advantageously injected *via* intravenous, intranasal (aerosolization), intramuscular, intradermal, and subcutaneous routes. A polynucleotide in a naked form can advantageously be administered *via* the intramuscular, intradermal, or subcutaneous routes.

Although not absolutely required, such a composition can also contain an adjuvant. If so, a systemic adjuvant that does not require concomitant administration in order to exhibit an adjuvant effect is preferable such as, *e.g.*, QS21, which is described in U.S. Patent 5,057,546.

The sequence information provided in the present application enables the design of specific nucleotide probes and primers that can be useful in diagnosis. Accordingly, in a fifth aspect of the invention, there is provided a nucleotide probe or primer having a sequence found in or derived by degeneracy of the genetic code from a sequence shown in SEQ ID NO: 1.

The term "probe" as used in the present application refers to DNA (preferably single stranded) or RNA molecules (or modifications or combinations thereof) that hybridize under the stringent conditions, as defined above, to nucleic acid molecules having sequences homologous to those shown in SEQ ID NOS: 1 and 2, or to a complementary or anti-sense sequence.

Generally, probes are significantly shorter than full-length sequences shown in SEQ ID NOS: 1 and 2; for example, they can contain from about 5 to about 100, preferably from about 10 to about 80 nucleotides. In particular, probes have sequences that are at least 75%, preferably at least 85%, more preferably 95% homologous to a portion of a sequence as shown in SEQ ID NOS: 1 and 2 or that are complementary to such sequences. Probes can contain modified bases

such as inosine, methyl-5-deoxycytidine, deoxyuridine, dimethylamino-5-deoxyuridine, or diamino-2, 6-purine. Sugar or phosphate residues can also be modified or substituted. For example, a deoxyribose residue can be replaced by a polyamide (Nielsen *et al.*, *Science* 254: 1497 (1991)) and phosphate residues can be replaced by ester groups such as diphosphate, alkyl, 5 arylphosphonate and phosphorothioate esters. In addition, the 2'-hydroxyl group on ribonucleotides can be modified by including, *e.g.*, alkyl groups.

Probes of the invention can be used in diagnostic tests, as capture or detection probes. Such capture probes can be conventionally immobilized on a solid support, directly or indirectly, by covalent means or by passive adsorption. A detection probe can be labelled by a detection 10 marker selected from radioactive isotopes; enzymes such as peroxidase, alkaline phosphatase, and enzymes able to hydrolyze a chromogenic, fluorogenic, or luminescent substrate; compounds that are chromogenic, fluorogenic, or luminescent; nucleotide base analogs; and biotin.

Probes of the invention can be used in any conventional hybridization technique, such as dot blot (Maniatis *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL* (1982) Cold Spring 15 Harbor Laboratory Press, Cold Spring Harbor, New York), Southern blot (*Southern, J. Mol. Biol.* 98: 503 (1975)), northern blot (identical to Southern blot to the exception that RNA is used as a target), or the sandwich technique (Dunn *et al.*, *Cell* 12: 23 (1977)). The latter technique involves the use of a specific capture probe and/or a specific detection probe with nucleotide sequences that at least partially differ from each other.

20 A primer is usually a probe of about 10 to about 40 nucleotides that is used to initiate enzymatic polymerization of DNA in an amplification process (*e.g.*, PCR), in an elongation process, or in a reverse transcription method. In a diagnostic method involving PCR, primers can be labelled.

Thus, the invention also encompasses (i) a reagent containing a probe of the invention for 25 detecting and/or identifying the presence of *Chlamydia* in a biological material; (ii) a method for detecting and/or identifying the presence of *Chlamydia* in a biological material, in which (a) a sample is recovered or derived from the biological material, (b) DNA or RNA is extracted from the material and denatured, and (c) exposed to a probe of the invention, for example, a capture, detection probe or both, under stringent hybridization conditions, such that hybridization is

detected; and (iii) a method for detecting and/or identifying the presence of *Chlamydia* in a biological material, in which (a) a sample is recovered or derived from the biological material, (b) DNA is extracted therefrom, (c) the extracted DNA is primed with at least one, and preferably two, primers of the invention and amplified by polymerase chain reaction, and (d) the amplified DNA fragment is produced.

As previously mentioned, polypeptides that can be produced upon expression of the newly identified open reading frames are useful vaccine agents.

Therefore, a sixth aspect of the invention features a substantially purified polypeptide or polypeptide derivative having an amino acid sequence encoded by a polynucleotide of the invention.

A "substantially purified polypeptide" is defined as a polypeptide that is separated from the environment in which it naturally occurs and/or that is free of the majority of the polypeptides that are present in the environment in which it was synthesized. For example, a substantially purified polypeptide is free from cytoplasmic polypeptides. Those skilled in the art will understand that the polypeptides of the invention can be purified from a natural source, *i.e.*, a *Chlamydia* strain, or can be produced by recombinant means.

Homologous polypeptides or polypeptide derivatives encoded by polynucleotides of the invention can be screened for specific antigenicity by testing cross-reactivity with an antiserum raised against the polypeptide of reference having an amino acid sequence as shown in SEQ ID NOS: 1 and 2. Briefly, a monospecific hyperimmune antiserum can be raised against a purified reference polypeptide as such or as a fusion polypeptide, for example, an expression product of MBP, GST, or His-tag systems or a synthetic peptide predicted to be antigenic. The homologous polypeptide or derivative screened for specific antigenicity can be produced as such or as a fusion polypeptide. In this latter case and if the antiserum is also raised against a fusion polypeptide, two different fusion systems are employed. Specific antigenicity can be determined according to a number of methods, including Western blot (Towbin *et al.*, *Proc. Natl. Acad. Sci. USA* 76: 4350 (1979)), dot blot, and ELISA, as described below.

In a Western blot assay, the product to be screened, either as a purified preparation or a total *E. coli* extract, is submitted to SDS-Page electrophoresis as described by Laemmli, *Nature* 227: 680 (1970). After transfer to a nitrocellulose membrane, the material is further incubated with the monospecific hyperimmune antiserum diluted in the range of dilutions from about 1:5 to about 1:5000, preferably from about 1:100 to about 1:500. Specific antigenicity is shown once a band corresponding to the product exhibits reactivity at any of the dilutions in the above range.

In an ELISA assay, the product to be screened is preferably used as the coating antigen. A purified preparation is preferred, although a whole cell extract can also be used. Briefly, about 100 µl of a preparation at about 10 µg protein/ml are distributed into wells of a 96-well polycarbonate ELISA plate. The plate is incubated for 2 hours at 37°C then overnight at 4°C. The plate is washed with phosphate buffer saline (PBS) containing 0.05% Tween 20 (PBS/Tween buffer). The wells are saturated with 250 µl PBS containing 1% bovine serum albumin (BSA) to prevent non-specific antibody binding. After 1 hour incubation at 37°C, the plate is washed with PBS/Tween buffer. The antiserum is serially diluted in PBS/Tween buffer containing 0.5% BSA. 100 µl of dilutions are added per well. The plate is incubated for 90 minutes at 37°C, washed and evaluated according to standard procedures. For example, a goat anti-rabbit peroxidase conjugate is added to the wells when specific antibodies were raised in rabbits. Incubation is carried out for 90 minutes at 37°C and the plate is washed. The reaction is developed with the appropriate substrate and the reaction is measured by colorimetry (absorbance measured spectrophotometrically). Under the above experimental conditions, a positive reaction is shown by O.D. values greater than a non immune control serum.

In a dot blot assay, a purified product is preferred, although a whole cell extract can also be used. Briefly, a solution of the product at about 100 µg/ml is serially two-fold diluted in 50 mM Tris-HCl (pH 7.5). 100 µl of each dilution are applied to a nitrocellulose membrane 0.45 µm set in a 96-well dot blot apparatus (Biorad). The buffer is removed by applying vacuum to the system. Wells are washed by addition of 50 mM Tris-HCl (pH 7.5) and the membrane is air-dried. The membrane is saturated in blocking buffer (50 mM Tris-HCl (pH 7.5) 0.15 M NaCl, 10 g/L skim milk) and incubated with an antiserum dilution from about 1:50 to about 1:5000, preferably about 1:500. The reaction is revealed according to standard procedures. For

example, a goat anti-rabbit peroxidase conjugate is added to the wells when rabbit antibodies are used. Incubation is carried out 90 minutes at 37°C and the blot is washed. The reaction is developed with the appropriate substrate and stopped. The reaction is measured visually by the appearance of a colored spot, *e.g.*, by colorimetry. Under the above experimental conditions, a positive reaction is shown once a colored spot is associated with a dilution of at least about 1:5, preferably of at least about 1:500.

Therapeutic or prophylactic efficacy of a polypeptide or derivative of the invention can be evaluated as described below.

According to a seventh aspect of the invention, there is provided (i) a composition of matter containing a polypeptide of the invention together with a diluent or carrier; in particular, (ii) a pharmaceutical composition containing a therapeutically or prophylactically effective amount of a polypeptide of the invention; (iii) a method for inducing an immune response against *Chlamydia* in a mammal, by administering to the mammal an immunogenically effective amount of a polypeptide of the invention to elicit an immune response, *e.g.*, a protective immune response to *Chlamydia*; and particularly, (iv) a method for preventing and/or treating a *Chlamydia* (*e.g.*, *C. trachomatis*, *C. psittaci*, *C. pneumoniae*, or *C. pecorum*) infection, by administering a prophylactic or therapeutic amount of a polypeptide of the invention to an individual in need. Additionally, the seventh aspect of the invention encompasses the use of a polypeptide of the invention in the preparation of a medicament for preventing and/or treating *Chlamydia* infection.

The immunogenic compositions of the invention can be administered by any conventional route in use in the vaccine field, in particular to a mucosal (*e.g.*, ocular, intranasal, pulmonary, oral, gastric, intestinal, rectal, vaginal, or urinary tract) surface or *via* the parenteral (*e.g.*, subcutaneous, intradermal, intramuscular, intravenous, or intraperitoneal) route. The choice of the administration route depends upon a number of parameters, such as the adjuvant associated with the polypeptide. For example, if a mucosal adjuvant is used, the intranasal or oral route will be preferred and if a lipid formulation or an aluminum compound is used, the parenteral route will be preferred. In the latter case, the subcutaneous or intramuscular route is most preferred. The choice can also depend upon the nature of the vaccine agent. For example,

a polypeptide of the invention fused to CTB or LTB will be best administered to a mucosal surface.

A composition of the invention can contain one or several polypeptides or derivatives of the invention. It can also contain at least one additional *Chlamydia* antigen, or a subunit,
5 fragment, homolog, mutant, or derivative thereof.

For use in a composition of the invention, a polypeptide or derivative thereof can be formulated into or with liposomes, preferably neutral or anionic liposomes, microspheres, ISCOMS, or virus-like-particles (VLPs) to facilitate delivery and/or enhance the immune response. These compounds are readily available to one skilled in the art; for example, see
10 LIPOSOMES: A PRACTICAL APPROACH (*supra*).

Adjuvants other than liposomes and the like can also be used and are known in the art. An appropriate selection can conventionally be made by those skilled in the art, for example, from the list provided below.

Administration can be achieved in a single dose or repeated as necessary at intervals as
15 can be determined by one skilled in the art. For example, a priming dose can be followed by three booster doses at weekly or monthly intervals. An appropriate dose depends on various parameters including the recipient (*e.g.*, adult or infant), the particular vaccine antigen, the route and frequency of administration, the presence/absence or type of adjuvant, and the desired effect (*e.g.*, protection and/or treatment), as can be determined by one skilled in the art. In general, a
0 vaccine antigen of the invention can be administered by a mucosal route in an amount from about 10 μ g to about 500 mg, preferably from about 1 mg to about 200 mg. For the parenteral route of administration, the dose usually should not exceed about 1 mg, preferably about 100 μ g.

When used as vaccine agents, polynucleotides and polypeptides of the invention can be used sequentially as part of a multistep immunization process. For example, a mammal can be
25 initially primed with a vaccine vector of the invention such as a pox virus, *e.g.*, *via* the parenteral route, and then boosted twice with the polypeptide encoded by the vaccine vector, *e.g.*, *via* the mucosal route. In another example, liposomes associated with a polypeptide or derivative of the

invention can also be used for priming, with boosting being carried out mucosally using a soluble polypeptide or derivative of the invention in combination with a mucosal adjuvant (*e.g.*, LT).

A polypeptide derivative of the invention is also useful as a diagnostic reagent for detecting the presence of anti-*Chlamydia* antibodies, *e.g.*, in a blood sample. Such polypeptides are about 5 to about 80, preferably about 10 to about 50 amino acids in length and can be labeled or unlabeled, depending upon the diagnostic method. Diagnostic methods involving such a reagent are described below.

Upon expression of a DNA molecule of the invention, a polypeptide or polypeptide derivative is produced and can be purified using known laboratory techniques. For example, the polypeptide or polypeptide derivative can be produced as a fusion protein containing a fused tail that facilitates purification. The fusion product can be used to immunize a small mammal, *e.g.*, a mouse or a rabbit, in order to raise antibodies against the polypeptide or polypeptide derivative (monospecific antibodies). The eighth aspect of the invention thus provides a monospecific antibody that binds to a polypeptide or polypeptide derivative of the invention.

By "monospecific antibody" is meant an antibody that is capable of reacting with a unique naturally-occurring *Chlamydia* polypeptide. An antibody of the invention can be polyclonal or monoclonal. Monospecific antibodies can be recombinant, *e.g.*, chimeric (*e.g.*, constituted by a variable region of murine origin associated with a human constant region), humanized (a human immunoglobulin constant backbone together with hypervariable region of animal, *e.g.*, murine, origin), and/or single chain. Both polyclonal and monospecific antibodies can also be in the form of immunoglobulin fragments, *e.g.*, F(ab)'2 or Fab fragments. The antibodies of the invention can be of any isotype, *e.g.*, IgG or IgA, and polyclonal antibodies can be of a single isotype or can contain a mixture of isotypes.

The antibodies of the invention, which are raised to a polypeptide or polypeptide derivative of the invention, can be produced and identified using standard immunological assays, *e.g.*, Western blot analysis, dot blot assay, or ELISA (see, *e.g.*, Coligan *et al.*, CURRENT PROTOCOLS IN IMMUNOLOGY (1994) John Wiley & Sons, Inc., New York, NY). The antibodies can be used in diagnostic methods to detect the presence of a *Chlamydia* antigen in a sample, such as a biological sample. The antibodies can also be used in affinity chromatography methods

for purifying a polypeptide or polypeptide derivative of the invention. As is discussed further below, such antibodies can be used in prophylactic and therapeutic passive immunization methods.

Accordingly, a ninth aspect of the invention provides (i) a reagent for detecting the presence of *Chlamydia* in a biological sample that contains an antibody, polypeptide, or polypeptide derivative of the invention; and (ii) a diagnostic method for detecting the presence of *Chlamydia* in a biological sample, by contacting the biological sample with an antibody, a polypeptide, or a polypeptide derivative of the invention, such that an immune complex is formed, and by detecting such complex to indicate the presence of *Chlamydia* in the sample or the organism from which the sample is derived.

Those skilled in the art will understand that the immune complex is formed between a component of the sample and the antibody, polypeptide, or polypeptide derivative, whichever is used, and that any unbound material can be removed prior to detecting the complex. As can be easily understood, a polypeptide reagent is useful for detecting the presence of anti-*Chlamydia* antibodies in a sample, *e.g.*, a blood sample, while an antibody of the invention can be used for screening a sample, such as a gastric extract or biopsy, for the presence of *Chlamydia* polypeptides.

For use in diagnostic applications, the reagent (*i.e.*, the antibody, polypeptide, or polypeptide derivative of the invention) can be in a free state or immobilized on a solid support, such as a tube, a bead, or any other conventional support used in the field. Immobilization can be achieved using direct or indirect means. Direct means include passive adsorption (non-covalent binding) or covalent binding between the support and the reagent. By "indirect means" is meant that an anti-reagent compound that interacts with a reagent is first attached to the solid support. For example, if a polypeptide reagent is used, an antibody that binds to it can serve as an anti-reagent, provided that it binds to an epitope that is not involved in the recognition of antibodies in biological samples. Indirect means can also employ a ligand-receptor system, for example, a molecule such as a vitamin can be grafted onto the polypeptide reagent and the corresponding receptor can be immobilized on the solid phase. This is illustrated by the biotin-streptavidin system. Alternatively, indirect means can be used, *e.g.*,

by adding to the reagent a peptide tail, chemically or by genetic engineering, and immobilizing the grafted or fused product by passive adsorption or covalent linkage of the peptide tail.

According to a tenth aspect of the invention, there is provided a process for purifying, from a biological sample, a polypeptide or polypeptide derivative of the invention, which involves carrying out antibody-based affinity chromatography with the biological sample, wherein the antibody is a monospecific antibody of the invention.

For use in a purification process of the invention, the antibody can be polyclonal or monospecific, and preferably is of the IgG type. Purified IgGs can be prepared from an antiserum using standard methods (see, *e.g.*, Coligan *et al.*, *supra*). Conventional chromatography supports, as well as standard methods for grafting antibodies, are disclosed in, *e.g.*, ANTIBODIES: A LABORATORY MANUAL, D. Lane, E. Harlow, Eds. (1988).

Briefly, a biological sample, such as an *C. pneumoniae* extract, preferably in a buffer solution, is applied to a chromatography material, preferably equilibrated with the buffer used to dilute the biological sample so that the polypeptide or polypeptide derivative of the invention (*i.e.*, the antigen) is allowed to adsorb onto the material. The chromatography material, such as a gel or a resin coupled to an antibody of the invention, can be in batch form or in a column. The unbound components are washed off and the antigen is then eluted with an appropriate elution buffer, such as a glycine buffer or a buffer containing a chaotropic agent, *e.g.*, guanidine HCl, or high salt concentration (*e.g.*, 3 M MgCl₂). Eluted fractions are recovered and the presence of the antigen is detected, *e.g.*, by measuring the absorbance at 280 nm.

An antibody of the invention can be screened for therapeutic efficacy as described as follows. According to an eleventh aspect of the invention, there is provided: (i) a composition of matter containing a monospecific antibody of the invention, together with a diluent or carrier; (ii) a pharmaceutical composition containing a therapeutically or prophylactically effective amount of a monospecific antibody of the invention, and (iii) a method for treating or preventing a *Chlamydia* (*e.g.*, *C. trachomatis*, *C. psittaci*, *C. pneumoniae* or *C. pecorum*) infection, by administering a therapeutic or prophylactic amount of a monospecific antibody of the invention to an individual in need. Additionally, the eleventh aspect of the invention encompasses the use

of a monospecific antibody of the invention in the preparation of a medicament for treating or preventing *Chlamydia* infection.

To this end, the monospecific antibody can be polyclonal or monoclonal, preferably of the IgA isotype (predominantly). In passive immunization, the antibody can be administered to a mucosal surface of a mammal, *e.g.*, the gastric mucosa, *e.g.*, orally or intragastrically, advantageously, in the presence of a bicarbonate buffer. Alternatively, systemic administration, not requiring a bicarbonate buffer, can be carried out. A monospecific antibody of the invention can be administered as a single active component or as a mixture with at least one monospecific antibody specific for a different *Chlamydia* polypeptide. The amount of antibody and the particular regimen used can be readily determined by one skilled in the art. For example, daily administration of about 100 to 1,000 mg of antibodies over one week, or three doses per day of about 100 to 1,000 mg of antibodies over two or three days, can be an effective regimens for most purposes.

Therapeutic or prophylactic efficacy can be evaluated using standard methods in the art, *e.g.*, by measuring induction of a mucosal immune response or induction of protective and/or therapeutic immunity, using, *e.g.*, the *C. pneumoniae* mouse model. Those skilled in the art will recognize that the *C. pneumoniae* strain of the model can be replaced with another *Chlamydia* strain. For example, the efficacy of DNA molecules and polypeptides from *C. pneumoniae* is preferably evaluated in a mouse model using an *C. pneumoniae* strain. Protection can be determined by comparing the degree of *Chlamydia* infection to that of a control group. Protection is shown when infection is reduced by comparison to the control group. Such an evaluation can be made for polynucleotides, vaccine vectors, polypeptides and derivatives thereof, as well as antibodies of the invention.

Adjuvants useful in any of the vaccine compositions described above are as follows.

Adjuvants for parenteral administration include aluminum compounds, such as aluminum hydroxide, aluminum phosphate, and aluminum hydroxy phosphate. The antigen can be precipitated with, or adsorbed onto, the aluminum compound according to standard protocols. Other adjuvants, such as RIBI (ImmunoChem, Hamilton, MT), can be used in parenteral administration.

Adjuvants for mucosal administration include bacterial toxins, *e.g.*, the cholera toxin (CT), the *E. coli* heat-labile toxin (LT), the *Clostridium difficile* toxin A and the *pertussis* toxin (PT), or combinations, subunits, toxoids, or mutants thereof. For example, a purified preparation of native cholera toxin subunit B (CTB) can be of use. Fragments, homologs, derivatives, and fusions to any of these toxins are also suitable, provided that they retain adjuvant activity. Preferably, a mutant having reduced toxicity is used. Suitable mutants are described, *e.g.*, in WO 95/17211 (Arg-7-Lys CT mutant), WO 96/6627 (Arg-192-Gly LT mutant), and WO 95/34323 (Arg-9-Lys and Glu-129-Gly PT mutant). Additional LT mutants that can be used in the methods and compositions of the invention include, *e.g.*, Ser-63-Lys, Ala-69-Gly, Glu-110-Asp, and Glu-112-Asp mutants. Other adjuvants, such as a bacterial monophosphoryl lipid A (MPLA) of, *e.g.*, *E. coli*, *Salmonella minnesota*, *Salmonella typhimurium*, or *Shigella flexneri*; saponins, or polylactide glycolide (PLGA) microspheres, can also be used in mucosal administration.

Adjuvants useful for both mucosal and parenteral administrations include polyphosphazene (WO 95/2415), DC-chol (3 b-(N-(N',N'-dimethyl aminomethane)-carbamoyl) cholesterol (U.S. Patent 5,283,185 and WO 96/14831) and QS-21 (WO 88/9336).

Any pharmaceutical composition of the invention, containing a polynucleotide, a polypeptide, a polypeptide derivative, or an antibody of the invention, can be manufactured in a conventional manner. In particular, it can be formulated with a pharmaceutically acceptable diluent or carrier, *e.g.*, water or a saline solution such as phosphate buffer saline. In general, a diluent or carrier can be selected on the basis of the mode and route of administration, and standard pharmaceutical practice. Suitable pharmaceutical carriers or diluents, as well as pharmaceutical necessities for their use in pharmaceutical formulations, are described in *Remington's Pharmaceutical Sciences*, a standard reference text in this field and in the USP/NF.

The invention also includes methods in which *Chlamydia* infection, are treated by oral administration of a *Chlamydia* polypeptide of the invention and a mucosal adjuvant, in combination with an antibiotic, an antacid, sucralfate, or a combination thereof. Examples of such compounds that can be administered with the vaccine antigen and the adjuvant are antibiotics, including, *e.g.*, macrolides, tetracyclines, and derivatives thereof (specific examples

of antibiotics that can be used include azithromycin or doxycyclin or immunomodulators such as cytokines or steroids. In addition, compounds containing more than one of the above-listed components coupled together, can be used. The invention also includes compositions for carrying out these methods, *i.e.*, compositions containing a *Chlamydia* antigen (or antigens) of the invention, an adjuvant, and one or more of the above-listed compounds, in a pharmaceutically acceptable carrier or diluent.

Amounts of the above-listed compounds used in the methods and compositions of the invention can readily be determined by one skilled in the art. In addition, one skilled in the art can readily design treatment/immunization schedules. For example, the non-vaccine components can be administered on days 1-14, and the vaccine antigen + adjuvant can be administered on days 7, 14, 21, and 28.

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples. These examples are described solely for purposes of illustration and are not intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may suggest or render expedient. Although specific terms have been employed herein, such terms are intended in a descriptive sense and not for purposes of limitation. Polypeptides having a sequence homologous to one of the sequences shown in SEQ ID NOS: 1 and 2, include naturally-occurring allelic variants, as well as mutants or any other non-naturally occurring variants that are analogous in terms of antigenicity, to a polypeptide.

As is known in the art, an allelic variant is an alternate form of a polypeptide that is characterized as having a substitution, deletion, or addition of one or more amino acids that does not alter the biological function of the polypeptide. By "biological function" is meant the function of the polypeptide in the cells in which it naturally occurs, even if the function is not necessary for the growth or survival of the cells. For example, the biological function of a porin is to allow the entry into cells of compounds present in the extracellular medium. The biological function is distinct from the antigenic function. A polypeptide can have more than one biological function.

Allelic variants are very common in nature. For example, a bacterial species, *e.g.*, *C. pneumoniae*, is usually represented by a variety of strains that differ from each other by minor allelic variations. Indeed, a polypeptide that fulfills the same biological function in different strains can have an amino acid sequence that is not identical in each of the strains. Such an
5 allelic variation may be equally reflected at the polynucleotide level.

Support for the use of allelic variants of polypeptide antigens comes from, *e.g.*, studies of the *Chlamydial* MOMP antigen. The amino acid sequence of the MOMP varies from strain to strain, yet cross-strain antibody binding plus neutralization of infectivity occurs, indicating that the MOMP, when used as an immunogen, is tolerant of amino acid variations.

10 Polynucleotides, *e.g.*, DNA molecules, encoding allelic variants can easily be retrieved by polymerase chain reaction (PCR) amplification of genomic bacterial DNA extracted by conventional methods. This involves the use of synthetic oligonucleotide primers matching upstream and downstream of the 5' and 3' ends of the encoding domain. Suitable primers can be designed according to the nucleotide sequence information provided in SEQ ID NOS: 1 and 2.
15 Typically, a primer can consist of 10 to 40, preferably 15 to 25 nucleotides. It may be also advantageous to select primers containing C and G nucleotides in a proportion sufficient to ensure efficient hybridization; *e.g.*, an amount of C and G nucleotides of at least 40%, preferably 50% of the total nucleotide amount.

Useful homologs that do not naturally occur can be designed using known methods for
20 identifying regions of an antigen that are likely to be tolerant of amino acid sequence changes and/or deletions. For example, sequences of the antigen from different species can be compared to identify conserved sequences.

Polypeptide derivatives that are encoded by polynucleotides of the invention include, *e.g.*, fragments, polypeptides having large internal deletions derived from full-length polypeptides,
25 and fusion proteins.

Polypeptide fragments of the invention can be derived from a polypeptide having a sequence homologous to any of the sequences shown in SEQ ID NO: 1, to the extent that the fragments retain the substantial antigenicity of the parent polypeptide (specific antigenicity).

Polypeptide derivatives can also be constructed by large internal deletions that remove a substantial part of the parent polypeptide, while retaining specific antigenicity. Generally, polypeptide derivatives should be about at least 12 amino acids in length to maintain antigenicity. Advantageously, they can be at least 20 amino acids, preferably at least 50 amino acids, more preferably at least 75 amino acids, and most preferably at least 100 amino acids in length.

Useful polypeptide derivatives, *e.g.*, polypeptide fragments, can be designed using computer-assisted analysis of amino acid sequences in order to identify sites in protein antigens having potential as surface-exposed, antigenic regions. *See e.g.*, Hughes *et al.*, *Infect. Immun.* 60(9):3497 1992.

Polypeptide fragments and polypeptides having large internal deletions can be used for revealing epitopes that are otherwise masked in the parent polypeptide and that may be of importance for inducing a protective T cell-dependent immune response. Deletions can also remove immunodominant regions of high variability among strains.

It is an accepted practice in the field of immunology to use fragments and variants of protein immunogens as vaccines, as all that is required to induce an immune response to a protein is a small (*e.g.*, 8 to 10 amino acid) immunogenic region of the protein. This has been done for a number of vaccines against pathogens other than *Chlamydia*. For example, short synthetic peptides corresponding to surface-exposed antigens of pathogens such as murine mammary tumor virus, peptide containing 11 amino acids; (*see e.g.*, Dion *et al.*, *Virology* 179:474-477 (1990)) Semliki Forest virus, peptide containing 16 amino acids (*see e.g.*, Snijders *et al.*, *J. Gen. Virol.* 72:557-565 (1991)), and canine parvovirus, 2 overlapping peptides, each containing 15 amino acids (*see e.g.*, Langeveld *et al.* *Vaccine* 12(15):1473-1480 (1994)), have been shown to be effective vaccine antigens against their respective pathogens.

Polynucleotides encoding polypeptide fragments and polypeptides having large internal deletions can be constructed using standard methods, for example, by PCR, including inverse PCR, by restriction enzyme treatment of the cloned DNA molecules, or by the method of Kunkel *et al.* (*Proc. Natl. Acad. Sci. USA* 82:448 (1985)) using biological material available at Stratagene.

A polypeptide derivative can also be produced as a fusion polypeptide that contains a polypeptide or a polypeptide derivative of the invention fused, *e.g.*, at the N- or C-terminal end, to any other polypeptide (hereinafter referred to as a peptide tail). Such a product can be easily obtained by translation of a genetic fusion, *i.e.*, a hybrid gene. Vectors for expressing fusion polypeptides are commercially available, such as the pMal-c2 or pMal-p2 systems of New England Biolabs, in which the peptide tail is a maltose binding protein, the glutathione-S-transferase system of Pharmacia, or the His-Tag system available from Novagen. These and other expression systems provide convenient means for further purification of polypeptides and derivatives of the invention.

Another particular example of fusion polypeptides included in invention includes a polypeptide or polypeptide derivative of the invention fused to a polypeptide having adjuvant activity, such as, *e.g.*, subunit B of either cholera toxin or *E. coli* heat-labile toxin. Several possibilities are can be used for achieving fusion. First, the polypeptide of the invention can be fused to the N-, or preferably, to the C-terminal end of the polypeptide having adjuvant activity. Second, a polypeptide fragment of the invention can be fused within the amino acid sequence of the polypeptide having adjuvant activity.

As stated above, the polynucleotides of the invention encode *Chlamydia* polypeptides in precursor or mature form. They can also encode hybrid precursors containing heterologous signal peptides, which can mature into polypeptides of the invention. By "heterologous signal peptide" is meant a signal peptide that is not found in the naturally-occurring precursor of a polypeptide of the invention.

A polynucleotide of the invention, having a homologous coding sequence, hybridizes, preferably under stringent conditions, to a polynucleotide having a sequence as shown in SEQ ID NOS: 1 or 2. Hybridization procedures are, *e.g.*, described in Ausubel *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons Inc. (1994), Silhavy *et al.* EXPERIMENTS WITH GENE FUSIONS, Cold Spring Harbor Laboratory Press (1984); Davis *et al.*, A MANUAL FOR GENETIC ENGINEERING: ADVANCED BACTERIAL GENETICS, Cold Spring Harbor Laboratory Press (1980). Important parameters that can be considered for optimizing hybridization conditions are reflected in a formula that allows calculation of a critical value, the

melting temperature above which two complementary DNA strands separate from each other. Casey and Davidson, *Nucl. Acid Res.* 4: 1539 (1977). This formula is as follows: $T_m = 81.5 + 0.41 \times (\% \text{ G+C}) + 16.6 \log (\text{cation ion concentration}) - 0.63 \times (\% \text{ formamide}) - 600/\text{base number}$. Under appropriate stringency conditions, hybridization temperature (T_h) is approximately
5 20-40°C, 20-25°C, or, preferably 30-40°C below the calculated T_m . Those skilled in the art will understand that optimal temperature and salt conditions can be readily determined empirically in preliminary experiments using conventional procedures.

For example, stringent conditions can be achieved, both for pre-hybridizing and hybridizing incubations, (i) within 4-16 hours at 42°C, in 6 x SSC containing 50% formamide or
10 (ii) within 4-16 hours at 65°C in an aqueous 6 x SSC solution (1 M NaCl, 0.1 M sodium citrate (pH 7.0)). Typically, hybridization experiments are performed at a temperature from 60 to 68°C, e.g., 65°C. At such a temperature, stringent hybridization conditions can be achieved in 6xSSC, preferably in 2xSSC or 1xSSC, more preferably in 0.5xSSC, 0.3xSSC or 0.1xSSC (in the absence of formamide). 1xSSC contains 0.15 M NaCl and 0.015 M sodium citrate.

15 For polynucleotides containing 30 to 600 nucleotides, the above formula is used and then is corrected by subtracting (600/polynucleotide size in base pairs). Stringency conditions are defined by a T_h that is 5 to 10°C below T_m .

Hybridization conditions with oligonucleotides shorter than 20-30 bases do not exactly follow the rules set forth above. In such cases, the formula for calculating the T_m is as follows:
20 $T_m = 4 \times (\text{G+C}) + 2 \times (\text{A+T})$. For example, an 18 nucleotide fragment of 50% G+C would have an approximate T_m of 54°C.

A polynucleotide molecule of the invention, containing RNA, DNA, or modifications or combinations thereof, can have various applications. For example, a DNA molecule can be used
25 (i) in a process for producing the encoded polypeptide in a recombinant host system, (ii) in the construction of vaccine vectors such as poxviruses, which are further used in methods and compositions for preventing and/or treating *Chlamydia* infection, (iii) as a vaccine agent (as well as an RNA molecule), in a naked form or formulated with a delivery vehicle and, (iv) in the construction of attenuated *Chlamydia* strains that can over-express a polynucleotide of the invention or express it in a non-toxic, mutated form.

According to a second aspect of the invention, there is therefore provided (i) an expression cassette containing a DNA molecule of the invention placed under the control of the elements required for expression, in particular under the control of an appropriate promoter; (ii) an expression vector containing an expression cassette of the invention; (iii) a prokaryotic or eukaryotic cell transformed or transfected with an expression cassette and/or vector of the invention, as well as (iv) a process for producing a polypeptide or polypeptide derivative encoded by a polynucleotide of the invention, which involves culturing a prokaryotic or eukaryotic cell transformed or transfected with an expression cassette and/or vector of the invention, under conditions that allow expression of the DNA molecule of the invention and, recovering the encoded polypeptide or polypeptide derivative from the cell culture.

A recombinant expression system can be selected from prokaryotic and eukaryotic hosts. Eukaryotic hosts include yeast cells (*e.g.*, *Saccharomyces cerevisiae* or *Pichia pastoris*), mammalian cells (*e.g.*, COS1, NIH3T3, or JEG3 cells), arthropods cells (*e.g.*, *Spodoptera frugiperda* (SF9) cells), and plant cells. Preferably, a prokaryotic host such as *E. coli* is used.

Bacterial and eukaryotic cells are available from a number of different sources to those skilled in the art, *e.g.*, the American Type Culture Collection (ATCC; Rockville, Maryland).

The choice of the expression system depends on the features desired for the expressed polypeptide. For example, it may be useful to produce a polypeptide of the invention in a particular lipidated form or any other form.

The choice of the expression cassette will depend on the host system selected as well as the features desired for the expressed polypeptide. Typically, an expression cassette includes a promoter that is functional in the selected host system and can be constitutive or inducible; a ribosome binding site; a start codon (ATG) if necessary, a region encoding a signal peptide, *e.g.*, a lipidation signal peptide; a DNA molecule of the invention; a stop codon; and optionally a 3' terminal region (translation and/or transcription terminator). The signal peptide encoding region is adjacent to the polynucleotide of the invention and placed in proper reading frame. The signal peptide-encoding region can be homologous or heterologous to the DNA molecule encoding the mature polypeptide and can be specific to the secretion apparatus of the host used for expression. The open reading frame constituted by the DNA molecule of the invention, solely or together

with the signal peptide, is placed under the control of the promoter so that transcription and translation occur in the host system. Promoters, signal peptide encoding regions are widely known and available to those skilled in the art and includes, for example, the promoter of *Salmonella typhimurium* (and derivatives) that is inducible by arabinose (promoter araB) and is functional in Gram-negative bacteria such as *E. coli* (as described in U.S. Patent No. 5,028,530 and in Cagnon *et al.*, *Protein Engineering* 4: 843 (1991); the promoter of the gene of bacteriophage T7 encoding RNA polymerase, that is functional in a number of *E. coli* strains expressing T7 polymerase (described in U.S. Patent No. 4,952,496); OspA lipidation signal peptide; and RlpB lipidation signal peptide. See Takase *et al.*, *J. Bact.* 169: 5692 (1987).

The expression cassette is typically part of an expression vector, which is selected for its ability to replicate in the chosen expression system. Expression vectors (*e.g.*, plasmids or viral vectors) can be chosen from those described in Pouwels *et al.* (Cloning Vectors: A Laboratory Manual 1985, Supp. 1987). They can be purchased from various commercial sources.

Methods for transforming/transfecting host cells with expression vectors will depend on the host system selected as described in Ausubel *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons Inc. (1994)).

Upon expression, a recombinant polypeptide of the invention (or a polypeptide derivative) is produced and remains in the intracellular compartment, is secreted/excreted in the extracellular medium or in the periplasmic space, or is embedded in the cellular membrane. The polypeptide can then be recovered in a substantially purified form from the cell extract or from the supernatant after centrifugation of the recombinant cell culture. Typically, the recombinant polypeptide can be purified by antibody-based affinity purification or by any other method that can be readily adapted by a person skilled in the art, such as by genetic fusion to a small affinity binding domain. Antibody-based affinity purification methods are also available for purifying a polypeptide of the invention extracted from a *Chlamydia* strain. Antibodies useful for purifying by immunoaffinity the polypeptides of the invention can be obtained as described below.

A polynucleotide of the invention can also be useful in the vaccine field, *e.g.*, for achieving DNA vaccination. There are two major possibilities, either using a viral or bacterial host as gene delivery vehicle (live vaccine vector) or administering the gene in a free form, *e.g.*,

inserted into a plasmid. Therapeutic or prophylactic efficacy of a polynucleotide of the invention can be evaluated as described below.

Accordingly, in a third aspect of the invention, there is provided (i) a vaccine vector such as a poxvirus, containing a DNA molecule of the invention, placed under the control of elements required for expression; (ii) a composition of matter containing a vaccine vector of the invention, together with a diluent or carrier; particularly, (iii) a pharmaceutical composition containing a therapeutically or prophylactically effective amount of a vaccine vector of the invention; (iv) a method for inducing an immune response against *Chlamydia* in a mammal (e.g., a human; alternatively, the method can be used in veterinary applications for treating or preventing *Chlamydia* infection of animals, e.g., cats or birds), which involves administering to the mammal an immunogenically effective amount of a vaccine vector of the invention to elicit an immune response, e.g., a protective or therapeutic immune response to *Chlamydia*; and particularly, (v) a method for preventing and/or treating a *Chlamydia* (e.g., *C. trachomatis*, *C. psittaci*, *C. pneumoniae*, *C. pecorum*) infection, which involves administering a prophylactic or therapeutic amount of a vaccine vector of the invention to an individual in need. Additionally, the third aspect of the invention encompasses the use of a vaccine vector of the invention in the preparation of a medicament for preventing and/or treating *Chlamydia* infection.

A vaccine vector of the invention can express one or several polypeptides or derivatives of the invention, as well as at least one additional *Chlamydia* antigen, fragment, homolog, mutant, or derivative thereof. In addition, it can express a cytokine, such as interleukin-2 (IL-2) or interleukin-12 (IL-12), that enhances the immune response (adjuvant effect). Thus, a vaccine vector can include an additional DNA sequence encoding, e.g., a chlamydial antigen, or a cytokine, placed under the control of elements required for expression in a mammalian cell.

Alternatively, a composition of the invention can include several vaccine vectors, each of them being capable of expressing a polypeptide or derivative of the invention. A composition can also contain a vaccine vector capable of expressing an additional *Chlamydia* antigen, or a subunit, fragment, homolog, mutant, or derivative thereof; or a cytokine such as IL-2 or IL-12.

In vaccination methods for treating or preventing infection in a mammal, a vaccine vector of the invention can be administered by any conventional route in use in the vaccine field,

particularly, to a mucosal (*e.g.*, ocular, intranasal, oral, gastric, pulmonary, intestinal, rectal, vaginal, or urinary tract) surface or *via* the parenteral (*e.g.*, subcutaneous, intradermal, intramuscular, intravenous, or intraperitoneal) route. Preferred routes depend upon the choice of the vaccine vector. The administration can be achieved in a single dose or repeated at intervals.

- 5 The appropriate dosage depends on various parameters understood by skilled artisans such as the vaccine vector itself, the route of administration or the condition of the mammal to be vaccinated (weight, age and the like).

Live vaccine vectors available in the art include viral vectors such as adenoviruses and poxviruses as well as bacterial vectors, *e.g.*, *Shigella*, *Salmonella*, *Vibrio cholerae*, *Lactobacillus*,
10 Bacille bilié de Calmette-Guérin (BCG), and *Streptococcus*.

- An example of an adenovirus vector, as well as a method for constructing an adenovirus vector capable of expressing a DNA molecule of the invention, are described in U.S. Patent No. 4,920,209. Poxvirus vectors that can be used include, *e.g.*, vaccinia and canary pox virus, described in U.S. Patent No. 4,722,848 and U.S. Patent No. 5,364,773, respectively. Also see,
5 *e.g.*, Tartaglia *et al.*, *Virology* 188: 217 (1992) for a description of a vaccinia virus vector; and Taylor *et al.*, *Vaccine* 13: 539 (1995) for a reference of a canary pox. Poxvirus vectors capable of expressing a polynucleotide of the invention can be obtained by homologous recombination as described in Kieny *et al.*, *Nature* 312: 163 (1984) so that the polynucleotide of the invention is inserted in the viral genome under appropriate conditions for expression in mammalian cells.
20 Generally, the dose of vaccine viral vector, for therapeutic or prophylactic use, can be of from about 1×10^4 to about 1×10^{11} , advantageously from about 1×10^7 to about 1×10^{10} , preferably of from about 1×10^7 to about 1×10^9 plaque-forming units per kilogram. Preferably, viral vectors are administered parenterally; for example, in 3 doses, 4 weeks apart. Those skilled in the art recognize that it is preferable to avoid adding a chemical adjuvant to a composition containing a
25 viral vector of the invention and thereby minimizing the immune response to the viral vector itself.

Non-toxicogenic *Vibrio cholerae* mutant strains that are useful as a live oral vaccine are described in Mekalanos *et al.*, *Nature* 306:551 (1983) and U.S. Patent No. 4,882,278 (strain in which a substantial amount of the coding sequence of each of the two *ctxA* alleles has been

deleted so that no functional *cholerae* toxin is produced); WO 92/11354 (strain in which the *irgA* locus is inactivated by mutation; this mutation can be combined in a single strain with *ctxA* mutations); and WO 94/1533 (deletion mutant lacking functional *ctxA* and *attRS1* DNA sequences). These strains can be genetically engineered to express heterologous antigens, as described in WO 94/19482. An effective vaccine dose of a *Vibrio cholerae* strain capable of expressing a polypeptide or polypeptide derivative encoded by a DNA molecule of the invention can contain, e.g., about 1×10^5 to about 1×10^9 , preferably about 1×10^6 to about 1×10^8 viable bacteria in an appropriate volume for the selected route of administration. Preferred routes of administration include all mucosal routes; most preferably, these vectors are administered intranasally or orally.

Attenuated *Salmonella typhimurium* strains, genetically engineered for recombinant expression of heterologous antigens or not, and their use as oral vaccines are described in Nakayama *et al.*, Bio/Technology 6:693 (1998) and WO 92/11361. Preferred routes of administration include all mucosal routes; most preferably, these vectors are administered intranasally or orally.

Others bacterial strains useful as vaccine vectors are described in High *et al.*, EMBO (1992) 11:1991 and Sizemore *et al.*, Science (1995) 270:299 (*Shigella flexneri*); Medaglini *et al.*, Proc. Natl. Acad. Sci. USA (1995) 92:6868 (*Streptococcus gordonii*); and Flynn, Cell. Mol. Biol. (1994) 40 (suppl. I):31, WO 88/6626, WO 90/0594, WO 91/13157, WO 92/1796, and WO 92/21376 (Bacille Calmette Guerin).

In bacterial vectors, polynucleotide of the invention can be inserted into the bacterial genome or can remain in a free state, carried on a plasmid.

An adjuvant can also be added to a composition containing a vaccine bacterial vector. A number of adjuvants are known to those skilled in the art. Preferred adjuvants can be selected from the list provided below.

According to a fourth aspect of the invention, there is also provided (i) a composition of matter containing a polynucleotide of the invention, together with a diluent or carrier; (ii) a pharmaceutical composition containing a therapeutically or prophylactically effective amount of

a polynucleotide of the invention; (iii) a method for inducing an immune response against *Chlamydia*, in a mammal, by administering to the mammal, an immunogenically effective amount of a polynucleotide of the invention to elicit an immune response, *e.g.*, a protective immune response to *Chlamydia*; and particularly, (iv) a method for preventing and/or treating a *Chlamydia* (*e.g.*, *C. trachomatis*, *C. psittaci*, *C. pneumoniae*, or *C. pecorum*) infection, by administering a prophylactic or therapeutic amount of a polynucleotide of the invention to an individual in need. Additionally, the fourth aspect of the invention encompasses the use of a polynucleotide of the invention in the preparation of a medicament for preventing and/or treating *Chlamydia* infection. The fourth aspect of the invention preferably includes the use of a DNA molecule placed under conditions for expression in a mammalian cell, *e.g.*, in a plasmid that is unable to replicate in mammalian cells and to substantially integrate in a mammalian genome.

Polynucleotides (DNA or RNA) of the invention can also be administered as such to a mammal for vaccine, *e.g.*, therapeutic or prophylactic, purpose. When a DNA molecule of the invention is used, it can be in the form of a plasmid that is unable to replicate in a mammalian cell and unable to integrate in the mammalian genome. Typically, a DNA molecule is placed under the control of a promoter suitable for expression in a mammalian cell. The promoter can function ubiquitously or tissue-specifically. Examples of non-tissue specific promoters include the early Cytomegalovirus (CMV) promoter (described in U.S. Patent No. 4,168,062) and the Rous Sarcoma Virus promoter (described in Norton & Coffin, *Molec. Cell Biol.* 5:281 (1985)). The desmin promoter (Li *et al.*, *Gene* 78: 243 (1989); Li & Paulin, *J. Biol. Chem.* 266: 6562 (1991); and Li & Paulin, *J. Biol. Chem.* 268: 10403 (1993)) is tissue-specific and drives expression in muscle cells. More generally, useful vectors are described, *i.a.*, WO 94/21797 and Hartikka *et al.*, *Human Gene Therapy* 7: 1205 (1996).

For DNA/RNA vaccination, the polynucleotide of the invention can encode a precursor or a mature form. When it encodes a precursor form, the precursor form can be homologous or heterologous. In the latter case, a eukaryotic leader sequence can be used, such as the leader sequence of the tissue-type plasminogen factor (tPA).

A composition of the invention can contain one or several polynucleotides of the invention. It can also contain at least one additional polynucleotide encoding another *Chlamydia*

antigen such as urease subunit A, B, or both; or a fragment, derivative, mutant, or analog thereof. A polynucleotide encoding a cytokine, such as interleukin-2 (IL-2) or interleukin-12 (IL-12), can also be added to the composition so that the immune response is enhanced. These additional polynucleotides are placed under appropriate control for expression. Advantageously, DNA molecules of the invention and/or additional DNA molecules to be included in the same composition, can be carried in the same plasmid.

Standard techniques of molecular biology for preparing and purifying polynucleotides can be used in the preparation of polynucleotide therapeutics of the invention. For use as a vaccine, a polynucleotide of the invention can be formulated according to various methods.

First, a polynucleotide can be used in a naked form, free of any delivery vehicles, such as anionic liposomes, cationic lipids, microparticles, *e.g.*, gold microparticles, precipitating agents, *e.g.*, calcium phosphate, or any other transfection-facilitating agent. In this case, the polynucleotide can be simply diluted in a physiologically acceptable solution, such as sterile saline or sterile buffered saline, with or without a carrier. When present, the carrier preferably is isotonic, hypotonic, or weakly hypertonic, and has a relatively low ionic strength, such as provided by a sucrose solution, *e.g.*, a solution containing 20% sucrose.

Alternatively, a polynucleotide can be associated with agents that assist in cellular uptake. It can be, *i.a.*, (i) complemented with a chemical agent that modifies the cellular permeability, such as bupivacaine (see, *e.g.*, WO 94/16737), (ii) encapsulated into liposomes, or (iii) associated with cationic lipids or silica, gold, or tungsten microparticles.

Anionic and neutral liposomes are well-known in the art (see, *e.g.*, Liposomes: A Practical Approach, RPC New Ed, IRL Press (1990), for a detailed description of methods for making liposomes) and are useful for delivering a large range of products, including polynucleotides.

Cationic lipids are also known in the art and are commonly used for gene delivery. Such lipids include LipofectinTM also known as DOTMA (N-[1-(2,3-dioleoyloxy)propyl]-N,N,N-trimethylammonium chloride), DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane), DDAB (dimethyldioctadecylammonium

bromide), DOGS (dioctadecylamidoglycyl spermine) and cholesterol derivatives such as DC-Chol (3 beta-(N-(N',N'-dimethyl aminomethane)-carbamoyl) cholesterol). A description of these cationic lipids can be found in EP 187,702, WO 90/11092, U.S. Patent No. 5,283,185, WO 91/15501, WO 95/26356, and U.S. Patent No. 5,527,928. Cationic lipids for gene delivery
5 are preferably used in association with a neutral lipid such as DOPE (dioleoyl phosphatidylethanolamine), as, for example, described in WO 90/11092.

Other transfection-facilitating compounds can be added to a formulation containing cationic liposomes. A number of them are described in, *e.g.*, WO 93/18759, WO 93/19768, WO 94/25608, and WO 95/2397. They include, *i.a.*, spermine derivatives useful for facilitating
10 the transport of DNA through the nuclear membrane (see, for example, WO 93/18759) and membrane-permeabilizing compounds such as GALA, Gramicidine S, and cationic bile salts (see, for example, WO 93/19768).

Gold or tungsten microparticles can also be used for gene delivery, as described in WO 91/359, WO 93/17706, and Tang *et al.* (Nature (1992) 356:152). In this case, the
15 microparticle-coated polynucleotides can be injected *via* intradermal or intraepidermal routes using a needleless injection device ("gene gun"), such as those described in U.S. Patent No. 4,945,050, U.S. Patent No. 5,015,580, and WO 94/24263.

The amount of DNA to be used in a vaccine recipient depends, *e.g.*, on the strength of the promoter used in the DNA construct, the immunogenicity of the expressed gene product, the
20 condition of the mammal intended for administration (*e.g.*, the weight, age, and general health of the mammal), the mode of administration, and the type of formulation. In general, a therapeutically or prophylactically effective dose from about 1 µg to about 1 mg, preferably, from about 10 µg to about 800 µg and, more preferably, from about 25 µg to about 250 µg, can be administered to human adults. The administration can be achieved in a single dose or
25 repeated at intervals.

The route of administration can be any conventional route used in the vaccine field. As general guidance, a polynucleotide of the invention can be administered *via* a mucosal surface, *e.g.*, an ocular, intranasal, pulmonary, oral, intestinal, rectal, vaginal, and urinary tract surface; or *via* a parenteral route, *e.g.*, by an intravenous, subcutaneous, intraperitoneal, intradermal,

intraepidermal, or intramuscular route. The choice of the administration route will depend on, *e.g.*, the formulation that is selected. A polynucleotide formulated in association with bupivacaine is advantageously administered into muscles. When a neutral or anionic liposome or a cationic lipid, such as DOTMA or DC-Chol, is used, the formulation can be advantageously injected *via* intravenous, intranasal (aerosolization), intramuscular, intradermal, and subcutaneous routes. A polynucleotide in a naked form can advantageously be administered *via* the intramuscular, intradermal, or sub-cutaneous routes.

Although not absolutely required, such a composition can also contain an adjuvant. If so, a systemic adjuvant that does not require concomitant administration in order to exhibit an adjuvant effect is preferable such as, *e.g.*, QS21, which is described in U.S. Patent No. 5,057,546.

The sequence information provided in the present application enables the design of specific nucleotide probes and primers that can be useful in diagnosis. Accordingly, in a fifth aspect of the invention, there is provided a nucleotide probe or primer having a sequence found in or derived by degeneracy of the genetic code from a sequence shown in SEQ ID NOS: 1 or 2.

The term "probe" as used in the present application refers to DNA (preferably single stranded) or RNA molecules (or modifications or combinations thereof) that hybridize under the stringent conditions, as defined above, to nucleic acid molecules having sequences homologous to those shown in SEQ ID NOS: 1 and 2, or to a complementary or anti-sense sequence. Generally, probes are significantly shorter than full-length sequences shown in SEQ ID NOS: 1 and 2; for example, they can contain from about 5 to about 100, preferably from about 10 to about 80 nucleotides. In particular, probes have sequences that are at least 75%, preferably at least 85%, more preferably 95% homologous to a portion of a sequence as shown in SEQ ID NOS: 1 and 2 or that are complementary to such sequences. Probes can contain modified bases such as inosine, methyl-5-deoxycytidine, deoxyuridine, dimethylamino-5-deoxyuridine, or diamino-2, 6-purine. Sugar or phosphate residues can also be modified or substituted. For example, a deoxyribose residue can be replaced by a polyamide (Nielsen *et al.*, Science (1991) 254:1497) and phosphate residues can be replaced by ester groups such as diphosphate, alkyl, arylphosphonate and phosphorothioate esters. In addition, the 2'-hydroxyl group on ribonucleotides can be modified by including, *e.g.*, alkyl groups.

Probes of the invention can be used in diagnostic tests, as capture or detection probes. Such capture probes can be conventionally immobilized on a solid support, directly or indirectly, by covalent means or by passive adsorption. A detection probe can be labeled by a detection marker selected from radioactive isotopes; enzymes such as peroxidase, alkaline phosphatase, and enzymes able to hydrolyze a chromogenic, fluorogenic, or luminescent substrate; compounds that are chromogenic, fluorogenic, or luminescent; nucleotide base analogs; and biotin.

Probes of the invention can be used in any conventional hybridization technique, such as dot blot (Maniatis *et al.*, Molecular Cloning: A Laboratory Manual (1982) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York), Southern blot (Southern, *J. Mol. Biol.* 98: 503 (1975)), northern blot (identical to Southern blot to the exception that RNA is used as a target), or the sandwich technique (Dunn *et al.*, *Cell* 12: 23 (1977)). The latter technique involves the use of a specific capture probe and/or a specific detection probe with nucleotide sequences that at least partially differ from each other.

A primer is usually a probe of about 10 to about 40 nucleotides that is used to initiate enzymatic polymerization of DNA in an amplification process (*e.g.*, PCR), in an elongation process, or in a reverse transcription method. In a diagnostic method involving PCR, primers can be labeled.

Thus, the invention also encompasses (i) a reagent containing a probe of the invention for detecting and/or identifying the presence of *Chlamydia* in a biological material; (ii) a method for detecting and/or identifying the presence of *Chlamydia* in a biological material, in which (a) a sample is recovered or derived from the biological material, (b) DNA or RNA is extracted from the material and denatured, and (c) exposed to a probe of the invention, for example, a capture, detection probe or both, under stringent hybridization conditions, such that hybridization is detected; and (iii) a method for detecting and/or identifying the presence of *Chlamydia* in a biological material, in which (a) a sample is recovered or derived from the biological material, (b) DNA is extracted therefrom, (c) the extracted DNA is primed with at least one, and preferably two, primers of the invention and amplified by polymerase chain reaction, and (d) the amplified DNA fragment is produced.

As previously mentioned, polypeptides that can be produced upon expression of the newly identified open reading frames are useful vaccine agents.

Therefore, a sixth aspect of the invention features a substantially purified polypeptide or polypeptide derivative having an amino acid sequence encoded by a polynucleotide of the invention.

A "substantially purified polypeptide" is defined as a polypeptide that is separated from the environment in which it naturally occurs and/or that is free of the majority of the polypeptides that are present in the environment in which it was synthesized. For example, a substantially purified polypeptide is free from cytoplasmic polypeptides. Those skilled in the art will understand that the polypeptides of the invention can be purified from a natural source, *i.e.*, a *Chlamydia* strain, or can be produced by recombinant means.

Homologous polypeptides or polypeptide derivatives encoded by polynucleotides of the invention can be screened for specific antigenicity by testing cross-reactivity with an antiserum raised against the polypeptide of reference having an amino acid sequence as shown in SEQ ID NO: 2. Briefly, a monospecific hyperimmune antiserum can be raised against a purified reference polypeptide as such or as a fusion polypeptide, for example, an expression product of MBP, GST, or His-tag systems or a synthetic peptide predicted to be antigenic. The homologous polypeptide or derivative screened for specific antigenicity can be produced as such or as a fusion polypeptide. In this latter case and if the antiserum is also raised against a fusion polypeptide, two different fusion systems are employed. Specific antigenicity can be determined according to a number of methods, including Western blot (Towbin *et al.*, *Proc. Natl. Acad. Sci. USA* 76: 4350 (1979)), dot blot, and ELISA, as described below.

In a Western blot assay, the product to be screened, either as a purified preparation or a total *E. coli* extract, is submitted to SDS-Page electrophoresis as described by Laemmli (*Nature* 227: 680 (1970)). After transfer to a nitrocellulose membrane, the material is further incubated with the monospecific hyperimmune antiserum diluted in the range of dilutions from about 1:5 to about 1:5000, preferably from about 1:100 to about 1:500. Specific antigenicity is shown once a band corresponding to the product exhibits reactivity at any of the dilutions in the above range.

In an ELISA assay, the product to be screened is preferably used as the coating antigen. A purified preparation is preferred, although a whole cell extract can also be used. Briefly, about 100 µl of a preparation at about 10 µg protein/ml are distributed into wells of a 96-well polycarbonate ELISA plate. The plate is incubated for 2 hours at 37°C then overnight at 4°C.

- 5 The plate is washed with phosphate buffer saline (PBS) containing 0.05% Tween 20 (PBS/Tween buffer). The wells are saturated with 250 µl PBS containing 1% bovine serum albumin (BSA) to prevent non-specific antibody binding. After 1 hour incubation at 37°C, the plate is washed with PBS/Tween buffer. The antiserum is serially diluted in PBS/Tween buffer containing 0.5% BSA. 100 µl of dilutions are added per well. The plate is incubated for
- 10 90 minutes at 37°C, washed and evaluated according to standard procedures. For example, a goat anti-rabbit peroxidase conjugate is added to the wells when specific antibodies were raised in rabbits. Incubation is carried out for 90 minutes at 37°C and the plate is washed. The reaction is developed with the appropriate substrate and the reaction is measured by colorimetry (absorbance measured spectrophotometrically). Under the above experimental conditions, a
- 15 positive reaction is shown by O.D. values greater than a non immune control serum.

- In a dot blot assay, a purified product is preferred, although a whole cell extract can also be used. Briefly, a solution of the product at about 100 µg/ml is serially two-fold diluted in 50 mM Tris-HCl (pH 7.5). 100 µl of each dilution are applied to a nitrocellulose membrane 0.45 µm set in a 96-well dot blot apparatus (Biorad). The buffer is removed by applying vacuum
- 20 to the system. Wells are washed by addition of 50 mM Tris-HCl (pH 7.5) and the membrane is air-dried. The membrane is saturated in blocking buffer (50 mM Tris-HCl (pH 7.5) 0.15 M NaCl, 10 g/L skim milk) and incubated with an antiserum dilution from about 1:50 to about 1:5000, preferably about 1:500. The reaction is revealed according to standard procedures. For
- 25 example, a goat anti-rabbit peroxidase conjugate is added to the wells when rabbit antibodies are used. Incubation is carried out 90 minutes at 37°C and the blot is washed. The reaction is developed with the appropriate substrate and stopped. The reaction is measured visually by the appearance of a colored spot, *e.g.*, by colorimetry. Under the above experimental conditions, a positive reaction is shown once a colored spot is associated with a dilution of at least about 1:5, preferably of at least about 1:500.

Therapeutic or prophylactic efficacy of a polypeptide or derivative of the invention can be evaluated as described below.

According to a seventh aspect of the invention, there is provided (i) a composition of matter containing a polypeptide of the invention together with a diluent or carrier; in particular, (ii) a pharmaceutical composition containing a therapeutically or prophylactically effective amount of a polypeptide of the invention; (iii) a method for inducing an immune response against *Chlamydia* in a mammal, by administering to the mammal an immunogenically effective amount of a polypeptide of the invention to elicit an immune response, *e.g.*, a protective immune response to *Chlamydia*; and particularly, (iv) a method for preventing and/or treating a *Chlamydia* (*e.g.*, *C. trachomatis*, *C. psittaci*, *C. pneumoniae*, or *C. pecorum*) infection, by administering a prophylactic or therapeutic amount of a polypeptide of the invention to an individual in need. Additionally, the seventh aspect of the invention encompasses the use of a polypeptide of the invention in the preparation of a medicament for preventing and/or treating *Chlamydia* infection.

The immunogenic compositions of the invention can be administered by any conventional route in use in the vaccine field, in particular to a mucosal (*e.g.*, ocular, intranasal, pulmonary, oral, gastric, intestinal, rectal, vaginal, or urinary tract) surface or *via* the parenteral (*e.g.*, subcutaneous, intradermal, intramuscular, intravenous, or intraperitoneal) route. The choice of the administration route depends upon a number of parameters, such as the adjuvant associated with the polypeptide. For example, if a mucosal adjuvant is used, the intranasal or oral route will be preferred and if a lipid formulation or an aluminum compound is used, the parenteral route will be preferred. In the latter case, the sub-cutaneous or intramuscular route is most preferred. The choice can also depend upon the nature of the vaccine agent. For example, a polypeptide of the invention fused to CTB or LTB will be best administered to a mucosal surface.

A composition of the invention can contain one or several polypeptides or derivatives of the invention. It can also contain at least one additional *Chlamydia* antigen, or a subunit, fragment, homolog, mutant, or derivative thereof.

For use in a composition of the invention, a polypeptide or derivative thereof can be formulated into or with liposomes, preferably neutral or anionic liposomes, microspheres, ISCOMS, or virus-like-particles (VLPs) to facilitate delivery and/or enhance the immune response. These compounds are readily available to one skilled in the art; for example, see

5 Liposomes: A Practical Approach (*supra*).

Adjuvants other than liposomes and the like can also be used and are known in the art. An appropriate selection can conventionally be made by those skilled in the art, for example, from the list provided below.

Administration can be achieved in a single dose or repeated as necessary at intervals as
10 can be determined by one skilled in the art. For example, a priming dose can be followed by three booster doses at weekly or monthly intervals. An appropriate dose depends on various parameters including the recipient (*e.g.*, adult or infant), the particular vaccine antigen, the route and frequency of administration, the presence/absence or type of adjuvant, and the desired effect (*e.g.*, protection and/or treatment), as can be determined by one skilled in the art. In general, a
15 vaccine antigen of the invention can be administered by a mucosal route in an amount from about 10 μ g to about 500 mg, preferably from about 1 mg to about 200 mg. For the parenteral route of administration, the dose usually should not exceed about 1 mg, preferably about 100 μ g.

When used as vaccine agents, polynucleotides and polypeptides of the invention can be used sequentially as part of a multistep immunization process. For example, a mammal can be
20 initially primed with a vaccine vector of the invention such as a pox virus, *e.g.*, *via* the parenteral route, and then boosted twice with the polypeptide encoded by the vaccine vector, *e.g.*, *via* the mucosal route. In another example, liposomes associated with a polypeptide or derivative of the invention can also be used for priming, with boosting being carried out mucosally using a soluble polypeptide or derivative of the invention in combination with a mucosal adjuvant (*e.g.*, LT).

25 A polypeptide derivative of the invention is also useful as a diagnostic reagent for detecting the presence of anti-*Chlamydia* antibodies, *e.g.*, in a blood sample. Such polypeptides are about 5 to about 80, preferably about 10 to about 50 amino acids in length and can be labeled or unlabeled, depending upon the diagnostic method. Diagnostic methods involving such a reagent are described below.

Upon expression of a DNA molecule of the invention, a polypeptide or polypeptide derivative is produced and can be purified using known laboratory techniques. For example, the polypeptide or polypeptide derivative can be produced as a fusion protein containing a fused tail that facilitates purification. The fusion product can be used to immunize a small mammal, *e.g.*, a mouse or a rabbit, in order to raise antibodies against the polypeptide or polypeptide derivative (monospecific antibodies). The eighth aspect of the invention thus provides a monospecific antibody that binds to a polypeptide or polypeptide derivative of the invention.

By "monospecific antibody" is meant an antibody that is capable of reacting with a unique naturally-occurring *Chlamydia* polypeptide. An antibody of the invention can be polyclonal or monoclonal. Monospecific antibodies can be recombinant, *e.g.*, chimeric (*e.g.*, constituted by a variable region of murine origin associated with a human constant region), humanized (a human immunoglobulin constant backbone together with hypervariable region of animal, *e.g.*, murine, origin), and/or single chain. Both polyclonal and monospecific antibodies can also be in the form of immunoglobulin fragments, *e.g.*, F(ab)'2 or Fab fragments. The antibodies of the invention can be of any isotype, *e.g.*, IgG or IgA, and polyclonal antibodies can be of a single isotype or can contain a mixture of isotypes.

The antibodies of the invention, which are raised to a polypeptide or polypeptide derivative of the invention, can be produced and identified using standard immunological assays, *e.g.*, Western blot analysis, dot blot assay, or ELISA (see, *e.g.*, Coligan *et al.*, Current Protocols in Immunology (1994) John Wiley & Sons, Inc., New York, NY). The antibodies can be used in diagnostic methods to detect the presence of a *Chlamydia* antigen in a sample, such as a biological sample. The antibodies can also be used in affinity chromatography methods for purifying a polypeptide or polypeptide derivative of the invention. As is discussed further below, such antibodies can be used in prophylactic and therapeutic passive immunization methods.

Accordingly, a ninth aspect of the invention provides (i) a reagent for detecting the presence of *Chlamydia* in a biological sample that contains an antibody, polypeptide, or polypeptide derivative of the invention; and (ii) a diagnostic method for detecting the presence of *Chlamydia* in a biological sample, by contacting the biological sample with an antibody, a

polypeptide, or a polypeptide derivative of the invention, such that an immune complex is formed, and by detecting such complex to indicate the presence of *Chlamydia* in the sample or the organism from which the sample is derived.

Those skilled in the art will understand that the immune complex is formed between a component of the sample and the antibody, polypeptide, or polypeptide derivative, whichever is used, and that any unbound material can be removed prior to detecting the complex. As can be easily understood, a polypeptide reagent is useful for detecting the presence of anti-*Chlamydia* antibodies in a sample, *e.g.*, a blood sample, while an antibody of the invention can be used for screening a sample, such as a gastric extract or biopsy, for the presence of *Chlamydia* polypeptides.

For use in diagnostic applications, the reagent (*i.e.*, the antibody, polypeptide, or polypeptide derivative of the invention) can be in a free state or immobilized on a solid support, such as a tube, a bead, or any other conventional support used in the field. Immobilization can be achieved using direct or indirect means. Direct means include passive adsorption (non-covalent binding) or covalent binding between the support and the reagent. By "indirect means" is meant that an anti-reagent compound that interacts with a reagent is first attached to the solid support. For example, if a polypeptide reagent is used, an antibody that binds to it can serve as an anti-reagent, provided that it binds to an epitope that is not involved in the recognition of antibodies in biological samples. Indirect means can also employ a ligand-receptor system, for example, a molecule such as a vitamin can be grafted onto the polypeptide reagent and the corresponding receptor can be immobilized on the solid phase. This is illustrated by the biotin-streptavidin system. Alternatively, indirect means can be used, *e.g.*, by adding to the reagent a peptide tail, chemically or by genetic engineering, and immobilizing the grafted or fused product by passive adsorption or covalent linkage of the peptide tail.

According to a tenth aspect of the invention, there is provided a process for purifying, from a biological sample, a polypeptide or polypeptide derivative of the invention, which involves carrying out antibody-based affinity chromatography with the biological sample, wherein the antibody is a monospecific antibody of the invention.

For use in a purification process of the invention, the antibody can be polyclonal or monospecific, and preferably is of the IgG type. Purified IgGs can be prepared from an antiserum using standard methods (see, *e.g.*, Coligan *et al.*, *supra*). Conventional chromatography supports, as well as standard methods for grafting antibodies, are disclosed in,
5 *e.g.*, Antibodies: A Laboratory Manual, D. Lane, E. Harlow, Eds. (1988).

Briefly, a biological sample, such as an *C. pneumoniae* extract, preferably in a buffer solution, is applied to a chromatography material, preferably equilibrated with the buffer used to dilute the biological sample so that the polypeptide or polypeptide derivative of the invention (*i.e.*, the antigen) is allowed to adsorb onto the material. The chromatography material, such as a
10 gel or a resin coupled to an antibody of the invention, can be in batch form or in a column. The unbound components are washed off and the antigen is then eluted with an appropriate elution buffer, such as a glycine buffer or a buffer containing a chaotropic agent, *e.g.*, guanidine HCl, or high salt concentration (*e.g.*, 3 M MgCl₂). Eluted fractions are recovered and the presence of the antigen is detected, *e.g.*, by measuring the absorbance at 280 nm.

An antibody of the invention can be screened for therapeutic efficacy as described as follows. According to an eleventh aspect of the invention, there is provided (i) a composition of matter containing a monospecific antibody of the invention, together with a diluent or carrier; (ii) a pharmaceutical composition containing a therapeutically or prophylactically effective amount of a monospecific antibody of the invention, and (iii) a method for treating or preventing a
20 *Chlamydia* (*e.g.*, *C. trachomatis*, *C. psittaci*, *C. pneumoniae* or *C. pecorum*) infection, by administering a therapeutic or prophylactic amount of a monospecific antibody of the invention to an individual in need. Additionally, the eleventh aspect of the invention encompasses the use of a monospecific antibody of the invention in the preparation of a medicament for treating or preventing *Chlamydia* infection.

To this end, the monospecific antibody can be polyclonal or monoclonal, preferably of the IgA isotype (predominantly). In passive immunization, the antibody can be administered to a mucosal surface of a mammal, *e.g.*, the gastric mucosa, *e.g.*, orally or intragastrically, advantageously, in the presence of a bicarbonate buffer. Alternatively, systemic administration, not requiring a bicarbonate buffer, can be carried out. A monospecific antibody of the invention
25

8. The polypeptide of claim 7, wherein the fusion polypeptide is a signal peptide.
9. The polypeptide of claim 7, wherein the fusion polypeptide comprises a heterologous polypeptide having adjuvant activity.
10. An expression cassette, comprising the polynucleotide of claim 1 operably linked to a promoter.
11. An expression vector, comprising the expression cassette of claim 10.
12. A host cell, comprising the expression cassette of claim 10.
13. The host cell of claim 10, wherein said host cell is a prokaryotic cell.
14. The host cell of claim 13, wherein said host cell is a eukaryotic cell.
15. A method for producing a recombinant polypeptide having SEQ ID NO: 2, comprising:
 - (a) culturing a host cell of claim 12, under conditions that allow the expression of the polypeptide; and
 - (b) recovering the recombinant polypeptide.
16. A vaccine vector, comprising the expression cassette of claim 10.
17. The vaccine vector of claim 16, wherein said host mammal is human.
18. The vaccine vector of claim 16, in a pharmaceutically acceptable excipient.
19. A pharmaceutical composition, comprising an immunologically effective amount of the vaccine vector of claim 14.

20. A method for inducing an immune response in a mammal, comprising:
administering to said mammal an immunologically effective amount of the vaccine vector of claim 16, wherein said administration induces an immune response.
21. A pharmaceutical composition, comprising an immunologically effective amount of the polypeptide of claim 5 and pharmaceutically acceptable diluent.
22. The pharmaceutical composition of claim 21, further comprising an adjuvant.
23. The pharmaceutical composition of claim 21, further comprising one or more known *Chlamydia* antigens.
24. A method for inducing an immune response in a mammal, comprising:
administering to said mammal an immunologically effective amount of the pharmaceutical composition of claim 21, wherein said administration induces an immune response.
25. A polynucleotide probe reagent capable of detecting the presence of *Chlamydia* in biological material, comprising a polynucleotide that hybridizes to the polynucleotide of claim 1 under stringent conditions.
26. The polynucleotide probe reagent of claim 25, wherein said reagent is a DNA primer.
27. A hybridization method for detecting the presence of *Chlamydia* in a sample, comprising the steps of:
- (a) obtaining polynucleotide from the sample;
 - (b) hybridizing said obtained polynucleotide with a polynucleotide probe reagent of claim 21 under conditions which allow for the hybridization of said probe and said sample; and
 - (c) detecting said hybridization of said detecting reagent with a polynucleotide in said sample.

28. An amplification method for detecting the presence of *Chlamydia* in a sample, comprising the steps of:
- (a) obtaining polynucleotide from the sample;
 - (c) amplifying said obtained polynucleotide using one or more polynucleotide probe reagents of claim 25; and
 - (d) detecting said amplified polypeptide.
29. A method for detecting the presence of *Chlamydia* in a sample comprising the steps of:
- (a) contacting said sample with a detecting reagent that binds to the polypeptide having SEQ ID NO: 2 to form a complex; and
 - (b) detecting said formed complex.
30. The method of claim 29, wherein said detecting reagent is an antibody.
31. The method of claim 30, wherein said antibody is a monoclonal antibody.
32. The method of claim 30, wherein said antibody is a polyclonal antibody.
33. An affinity chromatography method for substantially purifying a polypeptide having SEQ ID NO: 2, comprising the steps of:
- (a) contacting a sample containing said polypeptide with a detecting reagent that binds to said polypeptide to form a complex;
 - (c) isolating said formed complex;
 - (c) dissociating said formed complex; and
 - (d) isolating the dissociated polypeptide.
34. The method of claim 33, wherein said detecting reagent is an antibody.
35. The method of claim 34, wherein said antibody is a monoclonal antibody.

36. The method of claim 34, wherein said antibody is a polyclonal antibody.
37. An antibody that immunospecifically binds a polypeptide of claim 5, or a fragment or derivative of said antibody containing the binding domain thereof.

CHLAMYDIA ANTIGENS AND CORRESPONDING DNA FRAGMENTS AND USES THEREOF

ABSTRACT OF THE DISCLOSURE

In summary of this disclosure, the present invention provides a method of nucleic acid, including DNA, immunization of a host, including humans, against disease caused by infection by a strain of Chlamydia, specifically *C. pneumoniae*, employing a vector, containing a nucleotide sequence encoding an 98 kDa putative outer membrane protein of a strain of *Chlamydia pneumoniae* and a promoter to effect expression of the *98 kDa putative outer membrane protein* gene in the host. Modifications are possible within the scope of this invention.

TRADOCS-1257062.1(QXY#01!.DOC)

cgctcttacc tagtagaggt tgagtgaatt tcttgacttg tttctcctat tgggtgtatct 60
 cttaaaatat taaattcaaa atcaaagtat atattttaca atg aag tct tct ttc 115
 Met Lys Ser Ser Phe
 1 5

ccc aag ttt gta ttt tct aca ttt gct att ttc cct ttg tct atg att 163
 Pro Lys Phe Val Phe Ser Thr Phe Ala Ile Phe Pro Leu Ser Met Ile
 10 15 20

gct acc gag aca gtt ttg gat tca agt gcg agt ttc gat ggg aat aaa 211
 Ala Thr Glu Thr Val Leu Asp Ser Ser Ala Ser Phe Asp Gly Asn Lys
 25 30 35

aat ggt aat ttt tca gtt cgt gag agt cag gaa gat gct gga act acc 259
 Asn Gly Asn Phe Ser Val Arg Glu Ser Gln Glu Asp Ala Gly Thr Thr
 40 45 50

tac cta ttt aag gga aat gtc act cta gaa aat att cct gga aca ggc 307
 Tyr Leu Phe Lys Gly Asn Val Thr Leu Glu Asn Ile Pro Gly Thr Gly
 55 60 65

aca gca atc aca aaa agc tgt ttt aac aac act aag ggc gat ttg act 355
 Thr Ala Ile Thr Lys Ser Cys Phe Asn Asn Thr Lys Gly Asp Leu Thr
 70 75 80 85

FIG. 1A

ttc aca ggt aac ggg aac tct cta ttg ttc caa acg gtg gat gca ggg 403
 Phe Thr Gly Asn Gly Asn Ser Leu Leu Phe Gln Thr Val Asp Ala Gly
 90 95 100

act gta gca ggg gct gct gtt aac agc agc gtg gta gat aaa tct acc 451
 Thr Val Ala Gly Ala Ala Val Asn Ser Ser Val Val Asp Lys Ser Thr
 105 110 115

acg ttt ata ggg ttt tct tcg cta tct ttt att gcg tct cct gga agt 499
 Thr Phe Ile Gly Phe Ser Ser Leu Ser Phe Ile Ala Ser Pro Gly Ser
 120 125 130

tcg ata act acc ggc aaa gga gcc gtt agc tgc tct acg ggt agc ttg 547
 Ser Ile Thr Thr Gly Lys Gly Ala Val Ser Cys Ser Thr Gly Ser Leu
 135 140 145

agt ttg aca aaa aat gtc agt ttg ctc ttc agc aaa aac ttt tca acg 595
 Ser Leu Thr Lys Asn Val Ser Leu Leu Phe Ser Lys Asn Phe Ser Thr
 150 155 160 165

gat aat ggc ggt gct atc acc gca aaa act ctt tca tta aca ggg act 643
 Asp Asn Gly Gly Ala Ile Thr Ala Lys Thr Leu Ser Leu Thr Gly Thr
 170 175 180

aca atg tca gct ctg ttt tct gaa aat acc tcc tca aag aaa ggc gga 691
 Thr Met Ser Ala Leu Phe Ser Glu Asn Thr Ser Ser Lys Lys Gly Gly
 185 190 195

FIG. 1B

gcc att cag act tcc gat gcc ctt acc att act gga aac caa ggg gaa 739
 Ala Ile Gln Thr Ser Asp Ala Leu Thr Ile Thr Gly Asn Gln Gly Glu
 200 205 210

gtc tct ttt tct gac aat act tct tcg gat tct gga gct gca att ttt 787
 Val Ser Phe Ser Asp Asn Thr Ser Ser Asp Ser Gly Ala Ala Ile Phe
 215 220 225

aca gaa gcc tcg gtg act att tct aat aat gct aaa gtt tcc ttt att 835
 Thr Glu Ala Ser Val Thr Ile Ser Asn Asn Ala Lys Val Ser Phe Ile
 230 235 240 245

gac aat aag gtc aca gga gcg agc tcc tca aca acg ggg gat atg tca 883
 Asp Asn Lys Val Thr Gly Ala Ser Ser Ser Thr Thr Gly Asp Met Ser
 250 255 260

gga ggt gct atc tgt gct tat aaa act agt aca gat act aag gtc acc 931
 Gly Gly Ala Ile Cys Ala Tyr Lys Thr Ser Thr Asp Thr Lys Val Thr
 265 270 275

ctc act gga aat cag atg tta ctc ttc agc aac aat aca tcg aca aca 979
 Leu Thr Gly Asn Gln Met Leu Leu Phe Ser Asn Asn Thr Ser Thr Thr
 280 285 290

gcg gga gga gct atc tat gtg aaa aag ctc gaa ctg gct tcc gga gga 1027
 Ala Gly Gly Ala Ile Tyr Val Lys Lys Leu Glu Leu Ala Ser Gly Gly
 295 300 305

FIG. 1C

ctt	acc	cta	ttc	agt	aga	aat	agt	gtc	aat	gga	ggt	aca	gct	cct	aaa	1075
Leu	Thr	Leu	Phe	Ser	Arg	Asn	Ser	Val	Asn	Gly	Gly	Thr	Ala	Pro	Lys	
310					315					320					325	
ggt	gga	gcc	ata	gct	atc	gaa	gat	agt	ggg	gaa	ttg	agt	tta	tcc	gcc	1123
Gly	Gly	Ala	Ile	Ala	Ile	Glu	Asp	Ser	Gly	Glu	Leu	Ser	Leu	Ser	Ala	
				330					335					340		
gat	agt	ggt	gac	att	gtc	ttt	tta	ggg	aat	aca	gtc	act	tct	act	act	1171
Asp	Ser	Gly	Asp	Ile	Val	Phe	Leu	Gly	Asn	Thr	Val	Thr	Ser	Thr	Thr	
			345					350					355			
cct	ggg	acg	aat	aga	agt	agt	atc	gac	tta	gga	acg	agt	gca	aag	atg	1219
Pro	Gly	Thr	Asn	Arg	Ser	Ser	Ile	Asp	Leu	Gly	Thr	Ser	Ala	Lys	Met	
		360					365					370				
aca	gct	ttg	cgt	tct	gct	gct	ggt	aga	gcc	atc	tac	ttc	tat	gat	ccc	1267
Thr	Ala	Leu	Arg	Ser	Ala	Ala	Gly	Arg	Ala	Ile	Tyr	Phe	Tyr	Asp	Pro	
	375					380					385					
ata	act	aca	gga	tca	tcc	aca	aca	gtt	aca	gat	gtc	tta	aaa	gtt	aat	1315
Ile	Thr	Thr	Gly	Ser	Ser	Thr	Thr	Val	Thr	Asp	Val	Leu	Lys	Val	Asn	
390					395					400					405	
gag	act	ccg	gca	gat	tct	gca	cta	caa	tat	aca	ggg	aac	atc	atc	ttc	1363
Glu	Thr	Pro	Ala	Asp	Ser	Ala	Leu	Gln	Tyr	Thr	Gly	Asn	Ile	Ile	Phe	
				410					415				420			

FIG. 1D

aca gga gaa aag tta tca gag aca gag gcc gca gat tct aaa aat ctt	1411
Thr Gly Glu Lys Leu Ser Glu Thr Glu Ala Ala Asp Ser Lys Asn Leu	
425 430 435	
act tcg aag cta cta cag cct gta act ctt tca gga ggt act cta tct	1459
Thr Ser Lys Leu Leu Gln Pro Val Thr Leu Ser Gly Gly Thr Leu Ser	
440 445 450	
tta aaa cat gga gtg act ctg cag act cag gca ttc act caa cag gca	1507
Leu Lys His Gly Val Thr Leu Gln Thr Gln Ala Phe Thr Gln Gln Ala	
455 460 465	
gat tct cgt ctc gaa atg gac gta gga act act cta gaa cct gct gat	1555
Asp Ser Arg Leu Glu Met Asp Val Gly Thr Thr Leu Glu Pro Ala Asp	
470 475 480 485	
act agc acc ata aac aat ttg gtc att aac atc agt tct ata gac ggt	1603
Thr Ser Thr Ile Asn Asn Leu Val Ile Asn Ile Ser Ser Ile Asp Gly	
490 495 500	
gca aag aag gca aaa ata gaa acc aaa gct acg tca aaa aat ctg act	1651
Ala Lys Lys Ala Lys Ile Glu Thr Lys Ala Thr Ser Lys Asn Leu Thr	
505 510 515	
tta tct gga acc atc act tta ttg gac ccg acg ggc acg ttt tat gaa	1699
Leu Ser Gly Thr Ile Thr Leu Leu Asp Pro Thr Gly Thr Phe Tyr Glu	
520 525 530	

FIG. 1E

aat cat agt tta aga aat cct cag tcc tac gac atc tta gag ctc aaa	1747
Asn His Ser Leu Arg Asn Pro Gln Ser Tyr Asp Ile Leu Glu Leu Lys	
535 540 545	
gct tct gga act gta aca agc acc gca gtg act cca gat cct ata atg	1795
Ala Ser Gly Thr Val Thr Ser Thr Ala Val Thr Pro Asp Pro Ile Met	
550 555 560 565	
ggc gag aaa ttc cat tac ggc tat cag gga act tgg ggc cca att gtt	1843
Gly Glu Lys Phe His Tyr Gly Tyr Gln Gly Thr Trp Gly Pro Ile Val	
570 575 580	
tgg ggg aca ggg gct tct acg act gca acc ttc aac tgg act aaa act	1891
Trp Gly Thr Gly Ala Ser Thr Thr Ala Thr Phe Asn Trp Thr Lys Thr	
585 590 595	
ggc tat att cct aat ccc gag cgt atc ggc tct tta gtc cct aat agc	1939
Gly Tyr Ile Pro Asn Pro Glu Arg Ile Gly Ser Leu Val Pro Asn Ser	
600 605 610	
tta tgg aat gca ttt ata gat att agc tct ctc cat tat ctt atg gag	1987
Leu Trp Asn Ala Phe Ile Asp Ile Ser Ser Leu His Tyr Leu Met Glu	
615 620 625	
act gca aac gaa ggg ttg cag gga gac cgt gct ttt tgg tgt gct gga	2035
Thr Ala Asn Glu Gly Leu Gln Gly Asp Arg Ala Phe Trp Cys Ala Gly	
630 635 640 645	

FIG. 1F

tta tct aac ttc ttc cat aag gat agt aca aaa aca cga cgc ggg ttt	2083
Leu Ser Asn Phe Phe His Lys Asp Ser Thr Lys Thr Arg Arg Gly Phe	
650 655 660	
cgc cat ttg agt ggc ggt tat gtc ata gga gga aac cta cat act tgt	2131
Arg His Leu Ser Gly Gly Tyr Val Ile Gly Gly Asn Leu His Thr Cys	
665 670 675	
tca gat aag att ctt agt gct gca ttt tgt cag ctc ttt gga aga gat	2179
Ser Asp Lys Ile Leu Ser Ala Ala Phe Cys Gln Leu Phe Gly Arg Asp	
680 685 690	
aga gac tac ttt gta gct aag aat caa ggt aca gtc tac gga gga act	2227
Arg Asp Tyr Phe Val Ala Lys Asn Gln Gly Thr Val Tyr Gly Gly Thr	
695 700 705	
ctc tat tac cag cac aac gaa acc tat atc tct ctt cct tgc aaa cta	2275
Leu Tyr Tyr Gln His Asn Glu Thr Tyr Ile Ser Leu Pro Cys Lys Leu	
710 715 720 725	
cgg cct tgt tcg ttg tct tat gtt cct aca gag att cct gtt ctc ttt	2323
Arg Pro Cys Ser Leu Ser Tyr Val Pro Thr Glu Ile Pro Val Leu Phe	
730 735 740	
tca gga aac ctt agc tac acc cat acg gat aac gat ctg aaa acc aag	2371
Ser Gly Asn Leu Ser Tyr Thr His Thr Asp Asn Asp Leu Lys Thr Lys	
745 750 755	

FIG. 1G

tat aca aca tat cct act gtt aaa gga agc tgg ggg aat gat agt ttc	2419
Tyr Thr Thr Tyr Pro Thr Val Lys Gly Ser Trp Gly Asn Asp Ser Phe	
760 765 770	
gct tta gaa ttc ggt gga aga gct ccg att tgc tta gat gaa agt gct	2467
Ala Leu Glu Phe Gly Gly Arg Ala Pro Ile Cys Leu Asp Glu Ser Ala	
775 780 785	
cta ttt gag cag tac atg ccc ttc atg aaa ttg cag ttt gtc tat gca	2515
Leu Phe Glu Gln Tyr Met Pro Phe Met Lys Leu Gln Phe Val Tyr Ala	
790 795 800 805	
cat cag gaa ggt ttt aaa gaa cag gga aca gaa gct cgt gaa ttt gga	2563
His Gln Glu Gly Phe Lys Glu Gln Gly Thr Glu Ala Arg Glu Phe Gly	
810 815 820	
agt agc cgt ctt gtg aat ctt gcc tta cct atc ggg atc cga ttt gat	2611
Ser Ser Arg Leu Val Asn Leu Ala Leu Pro Ile Gly Ile Arg Phe Asp	
825 830 835	
aag gaa tca gac tgc caa gat gca acg tac aat cta act ctt ggt tat	2659
Lys Glu Ser Asp Cys Gln Asp Ala Thr Tyr Asn Leu Thr Leu Gly Tyr	
840 845 850	
act gtg gat ctt gtt cgt agt aac ccc gac tgt acg aca aca ctg cga	2707
Thr Val Asp Leu Val Arg Ser Asn Pro Asp Cys Thr Thr Thr Leu Arg	
855 860 865	

FIG. 1H

att agc ggt gat tct tgg aaa acc ttc ggt acg aat ttg gca aga caa	2755
Ile Ser Gly Asp Ser Trp Lys Thr Phe Gly Thr Asn Leu Ala Arg Gln	
870	875
880	885
gct tta gtc ctt cgt gca ggg aac cat ttt tgc ttt aac tca aat ttt	2803
Ala Leu Val Leu Arg Ala Gly Asn His Phe Cys Phe Asn Ser Asn Phe	
890	895
900	
gaa gcc ttt agc caa ttt tct ttt gaa ttg cgt ggg tca tct cgc aat	2851
Glu Ala Phe Ser Gln Phe Ser Phe Glu Leu Arg Gly Ser Ser Arg Asn	
905	910
915	
tac aat gta gac tta gga gca aaa tac caa ttc taa tgcgtagct	2897
Tyr Asn Val Asp Leu Gly Ala Lys Tyr Gln Phe	
920	925
ttggtaaaga gctccataca tcgaaggga aagagctttt aagatttctt gaaggctctt	2957
ttcgaatttcg atttccattt tagtgttttg ctaaaacact ttc	3000




FIG. 2B

BseNI	_____ ! ____ !! _____ *
BseRI	_____ ! ____ ! ____ ! _____
BsgI	_____ ! _____ ! ____
Bsh1236I	_____ ! _____
Bsh1365I	_____ ! _____
BsiBI	_____ ! _____
BsiCI	_____ ! _____
BsiHKA1	_____ ! _____ ! _____ !! _____ ! ____
BsiLI	_____ ! ____ ! _____ ! _____
BsiMI	_____ ! _____
BsiSI	_____ ! _____ ! _____ ! _____
BsiYI	! ____ * ____ ! _____ ! _____ !! _____
BsIZI	_____ ! ____ * _____
BslI	! ____ * ____ ! _____ ! _____ !! _____
BsmAI	____ ! ____ ! ____ ! _____ !! ____ ! _____ * ____ ! _____
BsmBI	_____ ! _____ ! _____
BsmFI	_____ ! ____ ! _____ ! _____ !! _____
BsmI	_____ ! _____ ! _____
Bsp119I	_____ ! _____
Bsp1286I	_____ ! _____ !! ____ ! _____ !! _____ ! ____
Bsp13I	_____ ! _____
BspEI	_____ ! _____
BspHI	_____ ! _____
BspLI	_____ ! ____ ! _____ ! _____ * ____ * _____ ! ____ ! _____
BspMI	_____ ! _____
BsrBRI	_____ ! _____
BsrFI	_____ ! _____
BsrI	_____ ! ____ !! _____ *

FIG. 2C

DpnI	_____*	_____!	_____!!	_____
DraI	_____!	_____!	_____!	_____
DraII	_____!	_____!	_____!	_____
DrdI	_____!	_____!	_____!	_____
DseDI	_____!	_____!	_____!	_____
Eam1104I	_____!	_____!	_____!	_____!
EarI	_____!	_____!	_____!	_____!
Eco130I	_____!	_____!	_____!	_____!
Eco24I	_____!	_____!	_____!	_____!
Eco31I	_____!	_____!	_____!	_____!
Eco47I	_____!	_____!	_____!	_____!
Eco57I	_____!	_____!	_____!	_____!
Eco88I	_____!	_____!	_____!	_____!
Eco91I	_____!	_____!	_____!	_____!
EcoNI	_____!!	_____!	_____!	_____!
EcoO65I	_____!	_____!	_____!	_____!
EcoRI	_____!	_____!	_____!	_____!
EcoRII	_____!	_____!	_____!	_____!
EcoT14I	_____!	_____!	_____!	_____!
EcoT22I	_____!	_____!	_____!	_____!
EcoT38I	_____!	_____!	_____!	_____!
ErhI	_____!	_____!	_____!	_____!
Esp1396I	_____!	_____!	_____!	_____!
Esp3I	_____!	_____!	_____!	_____!
FauI	_____!	_____!	_____!	_____!
Fnu4HI	_____*	_____!	_____!	_____!

FIG. 2E




FIG. 2F




FIG. 2G

PstI	_____!_____
RcaI	_____!_____
RsaI	_____!_!_____!_____!_!_____!_!_!_____
SacI	_____!_____!_____!_____!_!_____
SapI	_____!_____
Sau96I	_____!_*_____
ScrFI	_____!_!_____!_____
SduI	_____!_____!_!_____!_!_____!_!_____
SfaNI	_____!_!_____!_____!_____!_____
SfcI	_____!_____!_!_!_!_____!_____
SfuI	_____!_____
SpeI	_____!_____
Sse9I	!!_!_____!_____!_____!_!_____!!_!_!_!***!_____
SspI	_!_!_____
StyI	_____!_____
TaiI	_____!_____!_!_____!_____
TaqI	_____!_____!_!_!_!_!_!_____!_*
TfiI	_____!_____!_!_!_!_!_!_!_!_!_!_!_____
ThaI	_____!_____
Tru1I	_*_!_!_!_!_____!_*_!_!_!_!_!_!_!_!_!_!_!_____
Tru9I	_*_!_!_!_!_____!_*_!_!_!_!_!_!_!_!_!_!_!_____
TscI	_____!_____!_!_____!_____
TseI	_____*_!_____!_____!_____
Tsp45I	_____!_____!_!_!_*_____!_____
Tsp509I	!!_!_____!_____!_____!_!_____!!_!_!_!***!_____
TspEI	!!_!_____!_____!_____!_!_____!!_!_!_!***!_____
TspRI	_____!_____!_____!_____
Tth111I	_____!_____
Van91I	_____!_____
XbaI	_____!_____!_____
XcmI	_____!_!_____
XhoII	_____!_____!_!_____
XmnI	_____!_____!_____
Zsp2I	_____!_____

FIG. 2H




FIG. 3




FIG. 4

[illegible]

- ☒ I hereby claim the benefit under Title 35, United States Code, § 119(e) or §120 of any United States application(s), or §365(c) of any PCT International application(s) designating the United States of America listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

Application No. <i>(U.S.S.N.)</i>	Filing Date <i>(dd/mm/yy)</i>	Status <i>(Patented, Pending, Abandoned)</i>
60/106,070	29/10/98	Pending
60/122,066	01/03/99	Pending

PCT International Applications designative the United States:

PCT Appln No.	US Serial No.	PCT Filing Date	Status

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Attorney or Agent	Registration No.	Attorney or Agent	Registration No.
David B. Bernstein	32,112	A. Jason Mirabito	28,161
David F. Crosby	36,400	Michel Morency	Limited Recognition
James G. Cullem	43,569	John T. Prince	43,091
Brett N. Dorny	35,860	Brian Rosenbloom	41,276
Ivor R. Elrifi	39,529	Carol H. Peters	45,010
John A. Harre	37,345	Garfield B. Simms	45,109
David E. Johnson	41,874	Thomas M. Sullivan	39,392
Robert Klauszinski	42,742	Howard Susser	33,556
Kristin E. Konzak	P-44,848	Shelby J. Walker	45,192
Cynthia Kozakiewicz	42,764	Martin M. Zoltick	35,745
Leslie Meyer-Leon	37,381		
William Marino	44,219		

Address all telephone calls to Ivor R. Elrifi at telephone number 617/348-1747.
Address all correspondence to:

Ivor R. Elrifi, Ph.D.
Mintz, Levin, Cohn, Ferris, Glovsky and Popeo, P.C.
One Financial Center
Boston, Massachusetts 02111

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or patent issued thereon.

Inventor's Signature

Date

Full Name of Inventor: Andrew B. Murdin

Citizenship: Canada

Residence: Ontario, Canada

Post Office Address: 146 Rhodes Circle Newmarket, Ontario, Canada L3X 1V2

Inventor's Signature

Date

Full Name of Inventor: Raymond P. Oomen

Citizenship: Canada

Residence: Ontario, Canada

Post Office Address: RR No 1 Schomberg, Ontario, Canada L0G 1T0

Inventor's Signature

Date

Full Name of Inventor: Pamela L. Dunn

Citizenship: United Kingdom

Residence: Ontario, Canada

Post Office Address: 3700 Kaneff Cr., Apt. 703, Mississauga, Ontario, Canada L5A 4B8